



ISSN 0441-2516
NIRS-RSD-107

RADIOACTIVITY SURVEY DATA in Japan

Part 2
= Dietary Materials =

NUMBER 107
February 1996

National Institute of Radiological Sciences
Chiba, Japan

Radioactivity Survey Data
in Japan
Number 107

February 1996 part 1 = Dietary Materials =

Contents

	Page
Environmental and Dietary Materials	
(Japan Chemical Analysis Center)	
1. Collection and pretreatment of samples	1
2. Preparation of samples for analysis	3
3. Separation of Strontium-90 and Cesium-137	3
4. Determination of Stable Strontium, Calcium and Potassium	4
5. Counting	4
6. Results	5
(1) Strontium-90 and Cesium-137 in Total Diet	5
(2) Strontium-90 and Cesium-137 in Rice	9
(producing districts)	
(3)-1 Strontium-90 and Cesium-137 in Milk	10
(producing districts for domestic program)	
-2 Strontium-90 and Cesium-137 in Milk	11
(producing districts for WHO program)	
-3 Strontium-90 and Cesium-137 in Milk	12
(consuming districts)	
-4 Strontium-90 and Cesium-137 in Milk	14
(powderd milk)	
(4)-1 Strontium-90 and Cesium-137 in Vegetables	15
(producing districts)	
-2 Strontium-90 and Cesium-137 in Vegetables	17
(consuming districts)	
(5) Strontium-90 and Cesium-137 in Tea (Japanese Tea)	18
(6) Strontium-90 and Cesium-137 in Sea Fish	19
(7) Strontium-90 and Cesium-137 in Freshwater Fish	22
(8) Strontium-90 and Cesium-137 in Shellfish	24
(9) Strontium-90 and Cesium-137 in Seaweeds	26
7. Contents of Figure (Selected Locations)	28

Edited by National Institute of Radiological Sciences, under the supervision of Science and Technology Agency of Japanese Government.

Environmental and Dietary Materials*

(Japan Chemical Analysis Center)

1. Collection and pretreatment of samples

(1) Rain and dry fallout

Rain and dry fallout was collected monthly on a sampling tray, approximately 5000cm² in area, which was filled with water to a depth of 1 cm at the beginning of every month.

Strontium and cesium carrier solutions were added after the sample was filtered. The tray was washed with 5ℓ of distilled water and the washing was combined to the filtrate.

The sample was passed through a cation exchange column (500mℓ of Dowex 50W X8, 50~100 mesh, Na form) at a rate flow of 80mℓ/min.

(2) Airborne dust

Airborne dust was collected by an electrostatic precipitator or a filter air sampler for every three-months at a rate of more than 3000m³ per month.

The sampling was done 1 to 1.5 meters above the ground.

(3) Service water and freshwater

Service water, 100ℓ each, was collected at the intake of the water-treatment plant and at the tap after water was left running for five minutes. Strontium and cesium carriers were added to the filtered water sample. The subsequent process was the same as that described in the section (1). Freshwater was treated in the same way as the service water.

(4) Soil

Soil was collected from the location in the spacious and flat area without past surface disturbance caused by duststorms, inflow and outflow due to precipitation, etc.. Any places located under trees in a forest, in a stony area or inside of river banks were avoided. Soil was taken from two layers of different depths, 0-5cm and 5-20cm. The soil lumps were crushed by hands and dried in a drying oven regulated 105°C. The soil was then passed through a 2mm sieve to remove plant roots and pebbles.

(5) Sea water

Sea water was collected at the fixed stations

where the effect of terrestrial fresh water from rivers was expected to be negligibly small. A special consideration was also given to weather conditions.

The sampling was carried out when there was no rainfall for the last few days. To prevent contamination, water samples were collected at the bow of a sampling boat just before she stood still by scooping surface water using a polyethylene bucket.

Immediately after the collection, the samples were acidified to a pH lower than 3 by adding concentrated hydrochloric acid in a ratio of 1mℓ to 1ℓ of sea water, and then stored in 20ℓ polyethylene containers. The sampling equipments as well as containers were thoroughly rinsed with dilute hydrochloric acid and then with distilled water before use. Two hundred milliliters of sea water was also collected at the same stations for the determination of chlorinity.

(6) Sea sediments

Sediment was collected in the same area as that for the sea water sample, taking the following criteria into account:

- a. The depth of water exceeds 1m at low tide.
- b. No significant sedimental movement is observed in the vicinity of concern.
- c. Mud, silt and fine sand are preferable.

A conventional sediment sampling device was used for collecting the top few centimeters of surface sediment. Approximately 4kg of the sample in wet weight was spread on a stainless steel dish after removed of the pebbles, shells and other foreign materials, and dried in a drying oven regulated at 105°C.

(7) Total diet

A full one day ordinary diet including three meals, water, tea and other in-between snacks for five persons was collected as a sample of "total diet".

The sample in a large stainless steel pan was carbonized carefully by direct application of gas flame, and was transferred to a porcelain dish and then ashed at 450°C in an electric muffle furnace.

(8) Rice

Polished rice was collected in producing districts at the harvest and in consuming areas when new crops were first put on sale. The sample was carbonized and ashed in a porcelain dish.

* Samples were sent to the Center from 46 contracted prefectures.

(9) Milk

Raw milk was collected in producing districts and commercial milk was purchased in consuming districts. Milk in a stainless steel pan or a porcelain dish was evaporated to dryness followed by carbonization and ashing.

(10) Vegetables

Spinach and Japanese radish were selected as the representatives for leaf vegetables and for nonstarch roots, respectively. After removing soil, the edible part of vegetable sample was dried and carbonized in a stainless steel pan or a porcelain dish.

(11) Tea

Five hundred grams of manufactured green tea was collected, carbonized and ashed in a stainless steel pan or a porcelain dish.

(12) Fish, shellfish and seaweeds

a. Sea fish and freshwater fish

Fish was rinsed with water and blotted with a filter paper. Only the edible part was used in case of larger sized fish, and the whole part was used in case of smaller ones. Each sample was weighed and placed in a stainless steel pan or a porcelain dish. After carbonized, the sample was ashed in an electric muffle furnace.

b. Shellfish

Approximately 4kg of shellfish including the shells was collected or purchased. After removing the shells, it was treated in the same way as that for the sea fish.

c. Seaweeds

Edible seaweeds were collected and rinsed with water to remove sand and other adhering matters on the surface. These were removed of excess water, weighed dried and ashed.

Table 1 shows details of sample collection.

Table 1 Details of sample collection

Sample	Frequency of sampling	Quantity of sample
=Environmental materials=		
(1) Rain and dry fallout		
1. For domestic program	monthly	
2. For WHO program	monthly	
(2) Airborne dust	quarterly	>3000 m ³ /month
(3) Service water and freshwater		
1. Service water (source water)	semiyearly	100 ℥
2. Service water (tap water)	semiyearly	100 ℥
3. Freshwater	yearly (fishing season)	100 ℥
(4) Soil		
1. 0~ 5 cm	yearly	4 kg
2. 5~ 20cm	yearly	4 kg
(5) Sea water	yearly	40 ℥
(6) Sea sediments	yearly	4 kg
=Dietary materials=		
(7) Total diet	semiyearly	daily amount for 5 persons
(8) Rice		
1. Producing districts	yearly (harvesting season)	5 kg (polished rice)
2. Consuming districts	yearly (harvesting season)	5 kg (polished rice)
(9) Milk		
1. Producing districts for WHO program	quarterly (February, May, August and November)	3 ℥
2. Producing districts for domestic program	semiyearly (February and August)	3 ℥

Sample	Frequency of sampling	Quantity of sample
3. Consuming districts	semiyearly (February and August)	3 ℥
4. Powdered milk	semiyearly (April and October)	2~ 3 kg
(10) Vegetables		
1. Producing districts	yearly (harvesting season)	4 kg
2. Consuming districts	yearly (harvesting season)	4 kg
(11) Tea	yearly (the first harvesting season)	500g (manufactured tea)
(12) Fish, shellfish and seaweeds		
1. Sea fish	yearly (fishing season)	4 kg
2. Freshwater fish	yearly (fishing season)	4 kg
3. Shellfish	yearly (fishing season)	4 kg
4. Seaweeds	yearly (fishing season)	2~ 3 kg

2. Preparation of samples for analysis

(1) Rain, service water and freshwater

Strontium and cesium were eluted with hydrochloric acid from the cation exchange column. The residue of rain sample on the filter paper was ashed in an electric muffle furnace and the ash was dissolved in hydrochloric acid. The insoluble part was filtered and washed. The filtrate and the washings were combined to the previous eluate and used for radiochemical analysis.

(2) Soil and Sea sediment

Dried soil was crushed to smaller ones than 0.25 mm in size by a crusher. The sieved sample was ashed in an electric muffle furnace regulated at 450 °C. The sample was then heated with hydrochloric acid, strontium and cesium carrier solutions and the mixture was heated. The insoluble constituent was filtered off and washed with water.

The dried sample was crushed to smaller ones than 0.25mm by a crushing machine. The further preparation of the sample was the same as that described in the section 2-(2).

(3) Rice

The ashed sample was pulverized with a porcelain mortar and passed through a 0.35mm sieve. The sieved sample to which both strontium and cesium carriers were added, was digested with nitric acid by heating. After the sample was heated again with nitric acid to dryness, strontium and cesium were extracted with hydrochloric acid and water. The insoluble constituent was filtered and washed. The filtrate and washings were combined for subsequent radiochemical analysis.

(4) Airborne dust, diet, milk, vegetables, fish and shellfish, seaweeds, tea and others

These ashed samples were treated with the same

procedure as that described in the section 2-(4).

3. Separation of strontium-90 and cesium-137

(1) Strontium-90

Sample solutions, prepared as in the foregoing sections 2-(1) through 2-(4), were neutralized with sodium hydroxide. After sodium carbonate was added, the precipitate of strontium and calcium carbonates was separated. The supernatant solution was retained for cesium-137 determination. The carbonates were redissolved in hydrochloric acid and strontium and calcium were precipitated as oxalates. The precipitate was dissolved in nitric acid and strontium was separated from calcium by successive fuming nitric acid separation. Iron scavenging was made after addition of ferric iron carrier followed by barium chromate separation after addition of barium carrier to remove radium, its daughters and lead. Strontium was recovered as carbonate, and the precipitate was dried and weighed to determine strontium recovery. The strontium carbonate was dissolved in hydrochloric acid and iron carrier was added. The solution was allowed to stand for two weeks for strontium-90 and yttrium-90 to attain equilibrium. Yttrium-90 was coprecipitated with ferric hydroxide and the precipitate was filtered off, washed and counted.

(2) Cesium-137

The supernatant separated from the strontium fraction was acidified with hydrochloric acid. While stirring, cesium was adsorbed on the ammonium molybdate phosphate added.

After filtered off and washed with hydrochloric acid the precipitate was dissolved in 2.5N sodium hydroxide solution. The solution was adjusted to pH 8.2 with hydrochloric acid and allowed to cool. Resultant molybdenum hydroxide which separated out in the solution, was filtered off and washed with

water. EDTA was added to the filtrate and washings. Cesium and rubidium were adsorbed on a cation exchange column and cesium was separated from rubidium by eluting with hydrochloric acid.

The eluate was evaporated to dryness and was dissolved. The solution was filtered.

Chloroplatinic acid was added to precipitate cesium. The precipitate was filtered onto a tared paper using a demountable filter and washed with water and then ethanol. After drying, the chemical yield of cesium was determined by weighing the precipitate. Cesium-137 radioactivity was measured for this precipitate.

4. Determination of stable strontium, calcium and potassium

A weighed amount of soil or sea sediment was heated in a electric muffle furnace at 450°C and then treated with hydrochloric acid for extraction.

A weighed aliquot of ashed samples of total diet, vegetables, milk, fish, shellfish or seaweeds was

digested with hydrofluoric acid and nitric acid.

The extract was made up to an appropriate volume with dilute hydrochloric acid. The sample solution was analyzed for calcium by titration with standard potassium permanganate solution after separating calcium as oxalate. Atomic absorption spectroscopy was applied when appropriate. Stable strontium and potassium were determined by atomic absorption and flame emission spectrometry, respectively.

5. Counting

After the radiochemical separation the mounted precipitates were counted for activity using low background beta counters normally for 60 to 90 min. Net sample counting rates were corrected for counter efficiency, recovery, self-absorption and decay to obtain the content of strontium-90 and cesium-137 per sample aliquot. From the results, concentrations of these nuclides in the original samples were calculated.

6. Results

(1) Strontium-90 and Cesium-137 in Total Diet

(from May. 1993 to Sep. 1993)

-continued from No. 105 of this publication-

Table (1) Strontium-90 and Cesium-137 in Total Diet

Location	Ash	Ca	K	⁹⁰ Sr				¹³⁷ Cs			
	Ash(g/p·d)	Ca(mg/p·d)	K(mg/p·d)	(Bq/p·d)		(Bq/gCa)		(Bq/p·d)		(Bq/gK)	
May, 1993											
Yamagata, YAMAGATA	18.3	858	2250	0.045	± 0.015	0.052	± 0.018	0.027	± 0.0084	0.012	± 0.0037
Hamasaka-machi, HYOUGO	15.4	663	1960	0.056	± 0.0070	0.085	± 0.011	0.040	± 0.0082	0.020	± 0.0042
Kochi, KOCHI	13.2	509	1770	0.050	± 0.0087	0.098	± 0.017	0.055	± 0.0074	0.031	± 0.0042
Saga-machi, KOCHI	12.6	369	1460	0.11	± 0.014	0.29	± 0.039	0.041	± 0.0066	0.028	± 0.0045
June, 1993											
Sapporo, HOKKAIDOU	20.7	564	2440	0.12	± 0.017	0.21	± 0.030	0.073	± 0.0098	0.030	± 0.0040
Iwanai-machi, HOKKAIDOU	13.7	458	1680	0.044	± 0.012	0.096	± 0.027	0.034	± 0.0060	0.020	± 0.0036
Aomori, AOMORI	19.7	542	2260	0.065	± 0.013	0.12	± 0.024	0.027	± 0.0070	0.012	± 0.0031
Ajigasawa-machi, AOMORI	17.0	522	2210	0.12	± 0.017	0.22	± 0.033	0.038	± 0.0082	0.017	± 0.0037
Morioka, IWATE	16.4	559	1790	0.087	± 0.017	0.16	± 0.030	0.057	± 0.0099	0.032	± 0.0055
Iwaizumi-machi, IWATE	11.7	359	1310	0.054	± 0.011	0.15	± 0.030	0.044	± 0.0082	0.034	± 0.0063
Sagae, YAMAGATA	10.1	277	1330	0.088	± 0.0087	0.32	± 0.031	0.026	± 0.0077	0.019	± 0.0058
Fukushima, FUKUSHIMA	15.2	673	1760	0.035	± 0.011	0.052	± 0.016	0.032	± 0.0062	0.018	± 0.0035
Ookuma-machi, FUKUSHIMA	16.0	392	1730	0.053	± 0.011	0.14	± 0.028	0.033	± 0.0063	0.019	± 0.0036
Mito, IBARAKI	15.3	456	1890	0.057	± 0.0061	0.12	± 0.013	0.051	± 0.0063	0.027	± 0.0033
Tokai-mura, IBARAKI	17.1	628	2210	0.069	± 0.0068	0.11	± 0.011	0.054	± 0.0070	0.024	± 0.0032
Utsunomiya, TOCHIGI	11.1	319	1350	0.036	± 0.0076	0.11	± 0.024	0.041	± 0.0058	0.031	± 0.0043
Mooka, TOCHIGI	13.8	536	1860	0.047	± 0.011	0.088	± 0.020	0.046	± 0.0081	0.025	± 0.0043
Maebashi, GUNMA	13.2	440	1950	0.056	± 0.0066	0.13	± 0.015	0.027	± 0.0074	0.014	± 0.0038
Nakanojou-machi, GUNMA	14.4	530	1870	0.070	± 0.0078	0.13	± 0.015	0.054	± 0.0081	0.029	± 0.0043
Urawa, SAITAMA	16.8	860	1930	0.062	± 0.0081	0.072	± 0.0094	0.037	± 0.0070	0.019	± 0.0036
Kumagaya, SAITAMA	14.1	683	2170	0.045	± 0.012	0.066	± 0.018	0.047	± 0.0079	0.022	± 0.0036
Ichihara, CHIBA	15.5	676	1830	0.052	± 0.0091	0.077	± 0.014	0.037	± 0.0057	0.020	± 0.0031

Location	Ash	Ca	K	^{90}Sr				^{137}Cs			
	Ash(g/p·d)	Ca(mg/p·d)	K(mg/p·d)	(Bq/p·d)	(Bq/gCa)	(Bq/p·d)	(Bq/gK)	(Bq/p·d)	(Bq/gCa)	(Bq/p·d)	(Bq/gK)
Chikura-machi, CHIBA	16.9	428	2010	0.054 ± 0.0061	0.13 ± 0.014	0.027 ± 0.0056	0.013 ± 0.0028				
Shinjuku, TOKYO	13.4	451	1880	0.051 ± 0.0097	0.11 ± 0.022	0.042 ± 0.0063	0.022 ± 0.0034				
Hachijou-machi, TOKYO	16.0	1160	1690	0.050 ± 0.012	0.043 ± 0.011	0.037 ± 0.0071	0.022 ± 0.0042				
Yokohama, KANAGAWA	11.9	333	1860	0.035 ± 0.0052	0.11 ± 0.016	0.018 ± 0.0051	0.0099 ± 0.0027				
Hiratsuka, KANAGAWA	15.5	568	2370	0.042 ± 0.0060	0.075 ± 0.011	0.041 ± 0.0073	0.017 ± 0.0031				
Kashiwazaki, NIIGATA	18.8	422	2320	0.043 ± 0.011	0.10 ± 0.027	0.046 ± 0.0078	0.020 ± 0.0034				
Nishikawa-machi, NIIGATA	21.0	743	2770	0.048 ± 0.010	0.064 ± 0.014	0.084 ± 0.0095	0.030 ± 0.0035				
Toyama, TOYAMA	13.7	299	1750	0.067 ± 0.0061	0.23 ± 0.020	0.044 ± 0.0069	0.025 ± 0.0039				
Takaoka, TOYAMA	12.3	626	1540	0.056 ± 0.0074	0.090 ± 0.012	0.033 ± 0.0071	0.021 ± 0.0046				
Kanazawa, ISHIKAWA	15.8	502	1550	0.057 ± 0.0097	0.11 ± 0.019	0.055 ± 0.0068	0.035 ± 0.0044				
Yoshinodani-mura, ISHIKAWA	13.4	393	1670	0.044 ± 0.011	0.11 ± 0.027	0.12 ± 0.010	0.070 ± 0.0060				
Tsuruga, FUKUI	15.2	1070	1590	0.084 ± 0.013	0.079 ± 0.012	0.031 ± 0.0071	0.020 ± 0.0045				
Nagano, NAGANO	12.1	390	1440	0.048 ± 0.0064	0.12 ± 0.016	0.021 ± 0.0062	0.014 ± 0.0043				
Toubu-machi, NAGANO	15.2	507	2100	0.063 ± 0.011	0.12 ± 0.022	0.10 ± 0.009	0.048 ± 0.0043				
Gifu, GIFU	12.6	468	1750	0.043 ± 0.0064	0.092 ± 0.014	0.024 ± 0.0068	0.014 ± 0.0039				
Takayama, GIFU	15.7	555	2140	0.071 ± 0.013	0.13 ± 0.024	0.035 ± 0.0071	0.016 ± 0.0033				
Shizuoka, SHIZUOKA	15.0	708	2190	0.060 ± 0.013	0.085 ± 0.018	0.055 ± 0.0089	0.025 ± 0.0041				
Hamaoka-machi, SHIZUOKA	11.7	417	1820	0.041 ± 0.0049	0.099 ± 0.012	0.030 ± 0.0068	0.017 ± 0.0038				
Nagoya, AICHI	13.9	573	1790	0.050 ± 0.0061	0.087 ± 0.011	0.033 ± 0.0067	0.019 ± 0.0038				
Shinshiro, AICHI	17.4	946	2160	0.066 ± 0.0070	0.070 ± 0.0074	0.036 ± 0.0084	0.017 ± 0.0039				
Tsu, MIE	13.9	332	1580	0.058 ± 0.0065	0.18 ± 0.020	0.016 ± 0.0055	0.010 ± 0.0035				
Owase, MIE	15.7	512	1750	0.086 ± 0.0080	0.17 ± 0.016	0.030 ± 0.0069	0.017 ± 0.0040				
Otsu, SHIGA	17.3	536	2260	0.058 ± 0.0063	0.11 ± 0.012	0.068 ± 0.0079	0.030 ± 0.0035				
Imazu-machi, SHIGA	13.4	530	1920	0.088 ± 0.0074	0.17 ± 0.014	0.050 ± 0.0071	0.026 ± 0.0037				
Kyoto, KYOTO	17.1	715	2450	0.067 ± 0.0078	0.094 ± 0.011	0.056 ± 0.0084	0.023 ± 0.0035				

Location	Ash	Ca	K	^{90}Sr				^{137}Cs			
	Ash(g/p·d)	Ca(mg/p·d)	K(mg/p·d)	(Bq/p·d)	(Bq/gCa)	(Bq/p·d)	(Bq/gK)	(Bq/p·d)	(Bq/gCa)	(Bq/p·d)	(Bq/gK)
Maizuru, KYOTO	15.9	498	1850	0.091 ± 0.0087	0.18 ± 0.018	0.044 ± 0.0079	0.024 ± 0.0043				
Osaka, OSAKA	17.3	710	2400	0.071 ± 0.018	0.10 ± 0.025	0.041 ± 0.0093	0.017 ± 0.0039				
Sakai, OSAKA	12.4	503	1690	0.044 ± 0.0066	0.088 ± 0.013	0.041 ± 0.0084	0.024 ± 0.0050				
Kakogawa, HYOUGO	13.2	583	1960	0.066 ± 0.012	0.11 ± 0.020	0.040 ± 0.0066	0.021 ± 0.0034				
Kashiwara, NARA	14.5	698	1840	0.050 ± 0.0066	0.072 ± 0.0095	0.027 ± 0.0077	0.014 ± 0.0042				
Gojou, NARA	15.0	1100	2260	0.057 ± 0.012	0.052 ± 0.011	0.050 ± 0.0075	0.022 ± 0.0033				
Wakayama, WAKAYAMA	12.8	752	2000	0.041 ± 0.011	0.055 ± 0.014	0.023 ± 0.0051	0.012 ± 0.0026				
Tottori, TOTTORI	14.8	374	1940	0.073 ± 0.0078	0.20 ± 0.021	0.052 ± 0.0080	0.027 ± 0.0041				
Fukube-mura, TOTTORI	12.6	318	1860	0.057 ± 0.012	0.18 ± 0.036	0.042 ± 0.0060	0.023 ± 0.0032				
Matsue, SHIMANE	26.8	1240	4000	0.089 ± 0.013	0.072 ± 0.011	0.053 ± 0.0084	0.013 ± 0.0021				
Okayama, OKAYAMA	18.2	460	2620	0.087 ± 0.013	0.19 ± 0.028	0.037 ± 0.0067	0.014 ± 0.0026				
Kamisaibara-mura, OKAYAMA	17.0	599	2120	0.11 ± 0.017	0.19 ± 0.028	0.051 ± 0.0086	0.024 ± 0.0040				
Yamaguchi, YAMAGUCHI	12.9	439	1570	0.028 ± 0.010	0.064 ± 0.023	0.047 ± 0.0067	0.030 ± 0.0043				
Ajisu-machi, YAMAGUCHI	15.2	476	2460	0.054 ± 0.0063	0.11 ± 0.013	0.077 ± 0.0092	0.031 ± 0.0037				
Tokushima, TOKUSHIMA	16.7	653	2340	0.062 ± 0.0071	0.095 ± 0.011	0.034 ± 0.0068	0.015 ± 0.0029				
Takamatsu, KAGAWA	15.0	468	1860	0.036 ± 0.0065	0.077 ± 0.014	0.037 ± 0.0070	0.020 ± 0.0038				
Shirotori-machi, KAGAWA	15.5	391	1660	0.035 ± 0.0060	0.091 ± 0.015	0.029 ± 0.0065	0.017 ± 0.0039				
Matsuyama, EHIME	11.4	390	1410	0.029 ± 0.0098	0.074 ± 0.025	0.026 ± 0.0056	0.018 ± 0.0039				
Ikata-machi, EHIME	10.7	424	1280	0.019 ± 0.013	0.044 ± 0.030	0.020 ± 0.0069	0.016 ± 0.0054				
Fukuoka, FUKUOKA	9.01	201	874	0.021 ± 0.0047	0.10 ± 0.024	0.0048 ± 0.0053	0.0055 ± 0.0061				
Dazaifu, FUKUOKA	17.4	684	2330	0.055 ± 0.0089	0.081 ± 0.013	0.058 ± 0.0076	0.025 ± 0.0033				
Saga, SAGA	14.1	426	1590	0.026 ± 0.011	0.060 ± 0.026	0.021 ± 0.0057	0.013 ± 0.0036				
Nagasaki, NAGASAKI	14.8	577	1930	0.048 ± 0.0093	0.083 ± 0.016	0.042 ± 0.0071	0.022 ± 0.0037				
Matsuura, NAGASAKI	11.9	469	1440	0.039 ± 0.0093	0.083 ± 0.020	0.023 ± 0.0066	0.016 ± 0.0046				
Kumamoto, KUMAMOTO	14.8	439	1580	0.038 ± 0.0055	0.087 ± 0.013	0.032 ± 0.0067	0.020 ± 0.0042				

Location	Ash	Ca	K	^{90}Sr				^{137}Cs			
	Ash(g/p·d)	Ca(mg/p·d)	K(mg/p·d)	(Bq/p·d)		(Bq/gCa)		(Bq/p·d)		(Bq/gK)	
Aso-machi, KUMAMOTO	19.8	638	2540	0.096	\pm 0.0089	0.15	\pm 0.014	0.066	\pm 0.0091	0.026	\pm 0.0036
Ooita, OITA	15.6	539	1920	0.043	\pm 0.0050	0.080	\pm 0.0093	0.043	\pm 0.0064	0.022	\pm 0.0033
Saiki, OITA	12.3	442	1450	0.036	\pm 0.011	0.081	\pm 0.026	0.056	\pm 0.0084	0.039	\pm 0.0058
Miyazaki, MIYAZAKI	15.6	394	2050	0.052	\pm 0.0088	0.13	\pm 0.022	0.051	\pm 0.0065	0.025	\pm 0.0032
Takahara-machi, MIYAZAKI	21.9	698	3170	0.081	\pm 0.0078	0.12	\pm 0.011	0.17	\pm 0.012	0.053	\pm 0.0039
Kagoshima, KAGOSHIMA	12.2	428	1510	0.020	\pm 0.0090	0.046	\pm 0.021	0.034	\pm 0.0061	0.022	\pm 0.0041
Ookuchi, KAGOSHIMA	12.3	475	1510	0.035	\pm 0.010	0.074	\pm 0.021	0.034	\pm 0.0060	0.023	\pm 0.0040
Okinawa, Okinawa	14.7	1040	2060	0.058	\pm 0.012	0.055	\pm 0.011	0.031	\pm 0.0061	0.015	\pm 0.0029
July, 1993											
Ishinomaki, MIYAGI	18.2	931	2150	0.059	\pm 0.0061	0.063	\pm 0.0065	0.045	\pm 0.0068	0.021	\pm 0.0031
Onagawa-machi, MIYAGI	16.8	723	1930	0.059	\pm 0.0059	0.082	\pm 0.0082	0.044	\pm 0.0063	0.023	\pm 0.0033
Akita, AKITA	14.7	352	1880	0.052	\pm 0.011	0.15	\pm 0.032	0.029	\pm 0.0068	0.015	\pm 0.0036
Omagari, AKITA	14.5	450	1820	0.060	\pm 0.012	0.13	\pm 0.026	0.063	\pm 0.0078	0.035	\pm 0.0043
Fukui, FUKUI	17.2	637	2060	0.051	\pm 0.013	0.080	\pm 0.021	0.034	\pm 0.0074	0.016	\pm 0.0036
Koufu, YAMANASHI	13.4	331	1790	0.048	\pm 0.0067	0.15	\pm 0.020	0.030	\pm 0.0073	0.017	\pm 0.0041
Nirasaki, YAMANASHI	14.3	528	1750	0.083	\pm 0.0082	0.16	\pm 0.016	0.037	\pm 0.0073	0.021	\pm 0.0042
Kitayama-mura, WAKAYAMA	12.7	572	1530	0.072	\pm 0.0074	0.13	\pm 0.013	0.018	\pm 0.0062	0.012	\pm 0.0041
Kashima-machi, SHIMANE	18.6	1060	2310	0.081	\pm 0.012	0.076	\pm 0.011	0.040	\pm 0.0069	0.018	\pm 0.0030
Karatsu, SAGA	25.8	936	2550	0.081	\pm 0.014	0.086	\pm 0.015	0.027	\pm 0.0070	0.011	\pm 0.0028
Naha, Okinawa	15.0	453	2360	0.051	\pm 0.0053	0.11	\pm 0.012	0.046	\pm 0.0072	0.020	\pm 0.0031
August, 1993											
Hiroshima, HIROSHIMA	13.7	962	1610	0.039	\pm 0.010	0.040	\pm 0.010	0.0070	\pm 0.0062	0.0043	\pm 0.0039
Miyoshi, HIROSHIMA	12.3	328	1630	0.038	\pm 0.0049	0.12	\pm 0.015	0.031	\pm 0.0055	0.019	\pm 0.0034
September, 1993											
Kamiita-machi, TOKUSHIMA	15.3	594	1880	0.046	\pm 0.0060	0.077	\pm 0.010	0.036	\pm 0.0065	0.019	\pm 0.0034

(2) Strontium-90 and Cesium-137 in Rice (producing districts)

Sep. 1993

-continued from No. 105 of this publication-

Table (2) Strontium-90 and Cesium-137 in Rice

Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)	(Bq/gCa)		(Bq/kgwet)	(Bq/gK)	
September, 1993									
Gifu, GIFU	0.492	0.035	0.792	0.015 ± 0.0045	0.43 ± 0.13		0.0000 ± 0.0039	0.0000 ± 0.0049	
Matsusaka, MIE	0.618	0.026	1.14	0.0097 ± 0.0079	0.38 ± 0.31		0.0000 ± 0.0050	0.0000 ± 0.0044	
Koushi-machi, KUMAMOTO	0.473	0.016	1.06	0.0054 ± 0.0039	0.33 ± 0.24		0.0021 ± 0.0051	0.0019 ± 0.0048	
Sadohara-machi, MIYAZAKI	0.577	0.021	1.04	0.020 ± 0.0090	0.96 ± 0.43		0.0013 ± 0.0057	0.0013 ± 0.0055	

(3)-1 Strontium-90 and Cesium-137 in Milk (producing districts for domestic program)

(from Jun. 1993 to Aug. 1993)

-continued from No. 105 of this publication-

Table (3)-1 Strontium-90 and Cesium-137 in Milk

Location	Component			⁹⁰ Sr		¹³⁷ Cs		
	Ash(g/ℓ)	Ca(g/ℓ)	K(g/ℓ)	(Bq/ℓ)	(Bq/gCa)	(Bq/ℓ)	(Bq/gK)	
June, 1993								
Yamato-machi, SAGA	7.44	1.17	1.63	0.032 ± 0.0055	0.027 ± 0.0047	0.0048 ± 0.0047	0.0029 ± 0.0029	
August, 1993								
Aomori, AOMORI	7.18	1.04	1.61	0.16 ± 0.009	0.15 ± 0.009	0.18 ± 0.013	0.11 ± 0.008	
Takizawa-mura, IWATE	7.25	1.06	1.62	0.024 ± 0.0046	0.023 ± 0.0043	0.025 ± 0.0055	0.016 ± 0.0034	
Mito, IBARAKI	7.21	1.09	1.54	0.028 ± 0.0048	0.026 ± 0.0044	0.0054 ± 0.0049	0.0035 ± 0.0032	
Nishinasuno-machi, TOCHIGI	7.57	1.26	1.58	0.022 ± 0.0047	0.018 ± 0.0037	0.052 ± 0.0070	0.033 ± 0.0044	
Fujimi-mura, GUNMA	7.31	1.07	1.70	0.020 ± 0.0082	0.019 ± 0.0076	0.0065 ± 0.0041	0.0038 ± 0.0024	
Yachimata, CHIBA	7.65	1.17	1.69	0.027 ± 0.0049	0.023 ± 0.0042	0.045 ± 0.0066	0.026 ± 0.0039	
Tonami, TOYAMA	7.45	1.11	1.59	0.028 ± 0.0049	0.025 ± 0.0044	0.016 ± 0.0049	0.010 ± 0.0031	
Oshimizu-machi, ISHIKAWA	7.19	1.02	1.63	0.039 ± 0.0057	0.039 ± 0.0056	0.20 ± 0.013	0.12 ± 0.008	
Takane-machi, YAMANASHI	6.97	1.03	1.45	0.022 ± 0.0097	0.021 ± 0.0093	0.013 ± 0.0048	0.0093 ± 0.0033	
Kasamatsu-machi, GIFU	6.62	1.02	1.43	0.033 ± 0.0046	0.032 ± 0.0045	0.012 ± 0.0042	0.0082 ± 0.0029	
Oouchiyama-mura, MIE	7.21	1.06	1.60	0.028 ± 0.0087	0.026 ± 0.0082	0.0070 ± 0.0038	0.0044 ± 0.0024	
Hino-machi, SHIGA	7.24	1.13	1.59	0.020 ± 0.0042	0.018 ± 0.0037	0.013 ± 0.0044	0.0083 ± 0.0028	
Mihara-machi, HYOGO	6.84	1.06	1.53	0.026 ± 0.0096	0.024 ± 0.0091	0.0017 ± 0.0041	0.0011 ± 0.0027	
Oouda-machi, NARA	7.32	1.09	1.48	0.032 ± 0.0094	0.029 ± 0.0087	0.016 ± 0.0048	0.011 ± 0.0033	
Kamiita-machi, TOKUSHIMA	7.39	1.20	1.57	0.025 ± 0.0082	0.021 ± 0.0069	0.0004 ± 0.0042	0.0003 ± 0.0027	
Takase-machi, KAGAWA	7.50	1.11	1.60	0.014 ± 0.0093	0.013 ± 0.0084	0.0096 ± 0.0046	0.0060 ± 0.0029	
Matsuyama, EHIME	7.85	1.24	1.64	0.022 ± 0.0043	0.018 ± 0.0035	0.021 ± 0.0056	0.013 ± 0.0034	
Koushi-machi, KUMAMOTO	7.24	1.09	1.60	0.027 ± 0.0045	0.025 ± 0.0041	0.0000 ± 0.0041	0.0000 ± 0.0026	
Kujuu-machi, OITA	7.53	1.15	1.58	0.026 ± 0.0045	0.023 ± 0.0039	0.078 ± 0.0083	0.049 ± 0.0052	
Takahara-machi, MIYAZAKI	7.03	0.984	1.64	0.0059 ± 0.0036	0.0060 ± 0.0037	0.030 ± 0.0055	0.018 ± 0.0033	

(3)-2

Strontium-90 and Cesium-137 in Milk (producing districts for WHO program)

(from May, 1993 to Aug, 1993)

-continued from No. 105 of this publication-

Table (3)-2 Strontium-90 and Cesium-137 in Milk

Location	Component			⁹⁰ Sr				¹³⁷ Cs			
	Ash(g/ ℥)	Ca(g/ ℥)	K(g/ ℥)	(Bq/ ℥)	(Bq/gCa)		(Bq/ ℥)	(Bq/gK)			
May, 1993											
Hokudainoujou, HOKKAIDOU	7.36	1.10	1.66	0.035 ± 0.011	0.032	± 0.010	0.077 ± 0.0086	0.046 ± 0.0052			
Hachijo-Island, TOKYO	6.74	0.936	1.43	0.066 ± 0.011	0.071	± 0.012	0.077 ± 0.0086	0.054 ± 0.0060			
Nishikawa-machi, NIIGATA	7.30	1.03	1.69	0.022 ± 0.0047	0.021	± 0.0045	0.0085 ± 0.0050	0.0051 ± 0.0030			
Katsuyama, FUKUI	7.13	1.09	1.58	0.012 ± 0.0092	0.011	± 0.0085	0.043 ± 0.0071	0.027 ± 0.0045			
Nose-machi, OSAKA	7.40	1.12	1.56	0.053 ± 0.012	0.047	± 0.011	0.014 ± 0.0056	0.0091 ± 0.0036			
Hikawa-machi, SHIMANE	7.48	1.26	1.55	0.034 ± 0.0056	0.027	± 0.0045	0.074 ± 0.0087	0.048 ± 0.0056			
Takamiya-machi, HIROSHIMA	7.01	1.03	1.51	0.015 ± 0.0094	0.014	± 0.0091	0.0081 ± 0.0048	0.0054 ± 0.0032			
Kochi, KOCHI	7.31	1.10	1.48	0.057 ± 0.0067	0.052	± 0.0060	0.035 ± 0.0072	0.024 ± 0.0048			
Yasu-machi, FUKUOKA	7.14	1.06	1.52	0.026 ± 0.0098	0.025	± 0.0092	0.0056 ± 0.0048	0.0037 ± 0.0031			
Kajiki-machi, KAGOSHIMA	7.45	1.12	1.56	0.031 ± 0.0049	0.028	± 0.0043	0.029 ± 0.0066	0.019 ± 0.0042			
August, 1993											
Hokudainoujou, HOKKAIDOU	7.16	1.20	1.59	0.036 ± 0.0058	0.030	± 0.0048	0.11 ± 0.010	0.070 ± 0.0063			
Hachijo-Island, TOKYO	6.51	0.932	1.31	0.084 ± 0.0070	0.090	± 0.0075	0.079 ± 0.0086	0.060 ± 0.0066			
Nishikawa-machi, NIIGATA	7.36	1.07	1.72	0.020 ± 0.0080	0.018	± 0.0075	0.013 ± 0.0047	0.0077 ± 0.0027			
Katsuyama, FUKUI	7.30	1.04	1.59	0.021 ± 0.0052	0.021	± 0.0050	0.026 ± 0.0060	0.016 ± 0.0038			
Nose-machi, OSAKA	7.21	1.07	1.52	0.039 ± 0.0093	0.037	± 0.0087	0.0081 ± 0.0043	0.0053 ± 0.0028			
Hikawa-machi, SHIMANE	7.75	1.39	1.48	0.086 ± 0.012	0.062	± 0.0085	0.11 ± 0.010	0.072 ± 0.0068			
Takamiya-machi, HIROSHIMA	7.11	1.06	1.51	0.029 ± 0.0049	0.027	± 0.0047	0.0044 ± 0.0046	0.0029 ± 0.0030			
Kochi, KOCHI	7.32	1.13	1.57	0.085 ± 0.0085	0.076	± 0.0075	0.026 ± 0.0067	0.017 ± 0.0042			
Yasu-machi, FUKUOKA	6.82	1.04	1.46	0.034 ± 0.0060	0.032	± 0.0058	0.0075 ± 0.0064	0.0051 ± 0.0044			
Kajiki-machi, KAGOSHIMA	7.68	1.17	1.65	0.034 ± 0.0096	0.029	± 0.0082	0.021 ± 0.0062	0.013 ± 0.0038			

(3)-3 Strontium-90 and Cesium-137 in Milk (consuming districts)

(from May. 1993 to Sep. 1993)

-continued from No. 105 of this publication-

Table (3)-3 Strontium-90 and Cesium-137 in Milk

Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(g/ℓ)	Ca(g/ℓ)	K(g/ℓ)	(Bq/ℓ)	(Bq/gCa)	(Bq/ℓ)	(Bq/ℓ)	(Bq/gK)	
May, 1993									
Sendai, MIYAGI	7.18	1.09	1.54	0.029	± 0.0046	0.027	± 0.0042	0.019	± 0.0056
Hirosima, HIROSHIMA	7.04	1.06	1.54	0.036	± 0.0048	0.033	± 0.0045	0.034	± 0.0059
June, 1993									
Kyoto, KYOTO	7.22	1.07	1.59	0.026	± 0.0046	0.024	± 0.0043	0.014	± 0.0047
August, 1993									
Sapporo, HOKKAIDOU	7.32	1.14	1.58	0.077	± 0.012	0.068	± 0.010	0.055	± 0.0074
Akita, AKITA	6.41	0.917	1.37	0.033	± 0.0047	0.036	± 0.0051	0.050	± 0.0065
Yamagata, YAMAGATA	6.68	0.997	1.45	0.029	± 0.0048	0.029	± 0.0048	0.0061	± 0.0041
Fukushima, FUKUSHIMA	7.32	1.10	1.64	0.021	± 0.0088	0.019	± 0.0080	0.035	± 0.0060
Urawa, SAITAMA	7.04	1.06	1.53	0.014	± 0.0040	0.013	± 0.0038	0.015	± 0.0048
Shinjuku, TOKYO	7.33	1.05	1.54	0.029	± 0.0049	0.027	± 0.0047	0.041	± 0.0062
Yokohama, KANAGAWA	7.22	1.09	1.58	0.019	± 0.0045	0.018	± 0.0041	0.015	± 0.0052
Niigata, NIIGATA	7.57	1.13	1.64	0.023	± 0.0084	0.021	± 0.0075	0.018	± 0.0051
Fukui, FUKUI	7.29	1.09	1.60	0.029	± 0.0048	0.027	± 0.0044	0.016	± 0.0050
Nagano, NAGANO	6.10	0.926	1.36	0.016	± 0.0034	0.017	± 0.0037	0.0098	± 0.0036
Shizuoka, SHIZUOKA	7.14	1.07	1.55	0.019	± 0.0041	0.018	± 0.0039	0.020	± 0.0052
Nagoya, AICHI	7.39	1.08	1.56	0.029	± 0.0048	0.026	± 0.0044	0.0000	± 0.0039
Osaka, OSAKA	7.01	1.04	1.52	0.039	± 0.0058	0.038	± 0.0055	0.047	± 0.0075
Shinguu, WAKAYAMA	6.65	1.01	1.46	0.022	± 0.0077	0.022	± 0.0077	0.0089	± 0.0042
Yonago, TOTTORI	6.66	1.00	1.48	0.025	± 0.0080	0.025	± 0.0080	0.043	± 0.0066
Matsue, SHIMANE	7.21	1.10	1.55	0.015	± 0.0090	0.013	± 0.0082	0.0046	± 0.0046
Okayama, OKAYAMA	7.43	1.08	1.59	0.014	± 0.0039	0.013	± 0.0036	0.020	± 0.0053
Yamaguchi, YAMAGUCHI	7.02	1.04	1.51	0.030	± 0.0047	0.029	± 0.0045	0.017	± 0.0048
Matsuyama, EHIME	7.18	1.09	1.52	0.038	± 0.0050	0.035	± 0.0046	0.016	± 0.0053
Kochi, KOCHI	7.12	0.997	1.46	0.040	± 0.010	0.040	± 0.010	0.015	± 0.0045
Chikushino, FUKUOKA	6.86	1.04	1.48	0.025	± 0.0042	0.024	± 0.0040	0.017	± 0.0050

Location	Component			^{89}Sr			^{137}Cs		
	Ash(g/ℓ)	Ca(g/ℓ)	K(g/ℓ)	(Bq/ℓ)	(Bq/gCa)	(Bq/ℓ)	(Bq/gK)		
Nagasaki, NAGASAKI	6.91	1.06	1.49	0.027 ± 0.0044	0.026 ± 0.0042	0.016 ± 0.0044	0.010 ± 0.0029		
Kagoshima, KAGOSHIMA	7.21	1.08	1.51	0.011 ± 0.011	0.010 ± 0.011	0.021 ± 0.0057	0.014 ± 0.0038		
September, 1993									
Yonagusuku-mura, Okinawa	7.15	1.06	1.59	0.037 ± 0.0089	0.035 ± 0.0084	0.0035 ± 0.0036	0.0022 ± 0.0023		

(14)

(3)-4 Strontium-90 and Cesium-137 in Milk (powderd milk)

(Jul. 1993)

-continued from No. 105 of this publication-

Table (3)-4 Strontium-90 and Cesium-137 in Milk

Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kg)	(Bq/gCa)	(Bq/kg)	(Bq/kg)	(Bq/gK)	
July, 1993									
Sample A.	7.99	12.1	17.3	0.38 ± 0.015	0.032 ± 0.0012	0.36 ± 0.019	0.021 ± 0.0011		
Sample B.	2.50	3.35	5.43	0.042 ± 0.0064	0.013 ± 0.0019	0.057 ± 0.0085	0.011 ± 0.0016		
Sample C.	9.51	14.3	21.3	0.95 ± 0.026	0.067 ± 0.0018	2.3 ± 0.05	0.11 ± 0.002		
Sample D.	3.82	3.79	5.81	0.036 ± 0.0056	0.0095 ± 0.0015	0.054 ± 0.0079	0.0092 ± 0.0014		
Sample E.	2.42	3.63	5.61	0.086 ± 0.0079	0.024 ± 0.0022	0.20 ± 0.014	0.035 ± 0.0024		
Sample F.	2.51	3.46	5.20	0.069 ± 0.0076	0.020 ± 0.0022	0.27 ± 0.016	0.052 ± 0.0030		

* Skin milk

(4)-1 Strontium-90 and cesium-137 in Vegetables (producing districts)
(from Feb. 1993 to Aug. 1993)

-continued from No. 105 of this publication-

Table (4)-1 :Strontium-90 and cesium-137 in Vegetables

Location	Component			⁹⁰Sr			¹³⁷Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kg wet)	(Bq/g Ca)	(Bq/kg wet)	(Bq/g Ca)	(Bq/g K)	
(Cabbage)									
February, 1993									
Kumatori-machi, OSAKA (Chinese cabbage)	0.685	0.248	2.93	0.027	± 0.0040	0.11	± 0.016	0.0043	± 0.0035
June, 1993									
Oota, SHIMANE (Japanese radish)	0.901	1.14	2.67	1.9	± 0.05	1.7	± 0.04	2.5	± 0.04
February, 1993									
Hiroshima, HIROSHIMA	0.493	0.184	1.94	0.015	± 0.0050	0.083	± 0.027	0.0051	± 0.0051
May, 1993									
Tahara-machi, AICHI	0.592	0.128	2.63	0.022	± 0.0056	0.17	± 0.044	0.0000	± 0.0072
June, 1993									
Oota, SHIMANE	0.549	0.144	2.17	0.22	± 0.020	1.5	± 0.14	0.33	± 0.018
Koushi-machi, KUMAMOTO	0.528	0.186	2.10	0.044	± 0.014	0.23	± 0.073	0.018	± 0.0078
August, 1993									
Ishikari-machi, HOKKAIDO (Onion)	0.578	0.145	2.52	0.23	± 0.023	1.6	± 0.16	0.0056	± 0.0070
July, 1993									
Kumatori-machi, OSAKA (Potato)	0.360	0.149	1.46	0.027	± 0.0044	0.18	± 0.029	0.0034	± 0.0051
July, 1993									
Mutsu, AOMORI (Spinach)	0.838	0.019	3.85	0.019	± 0.0039	0.98	± 0.20	0.085	± 0.0080
February, 1993									
Hiroshima, HIROSHIMA	1.54	0.547	6.24	0.049	± 0.0053	0.089	± 0.0097	0.0038	± 0.0038
May, 1993									

(16)

Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)	(Bq/gCa)		(Bq/kgwet)	(Bq/gK)	
Tahara-machi, AICHI	1.60	0.325	7.08	0.023 ± 0.0052	0.071 ± 0.016		0.0045 ± 0.0070	0.00063 ± 0.00099	
Koushi-machi, KUMAMOTO	1.56	0.710	6.57	0.084 ± 0.015	0.12 ± 0.021		0.0040 ± 0.0072	0.0006 ± 0.0011	
August, 1993									
Ishikari-machi, HOKKAIDOU	2.17	0.675	9.37	0.15 ± 0.017	0.23 ± 0.026		0.015 ± 0.0074	0.0016 ± 0.00079	

(4)-2 Strontium-90 and cesium-137 in Vegetables (consuming districts)
 (from May, 1993 to Sep. 1993)

-continued from No. 105 of this publication-

Table (4)-2 :Strontium-90 and cesium-137 in Vegetables

Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)		(Bq/gCa)	(Bq/kgwet)		(Bq/gK)
(Japanese radish)									
September, 1993									
Urawa, SAITAMA (Spinach)	0.609	0.187	2.62	0.11	± 0.016	0.61	± 0.086	0.016	± 0.0057
May, 1993									
Sendai, MIYAGI	1.50	0.670	5.94	0.11	± 0.008	0.16	± 0.012	0.017	± 0.0078
June, 1993									
Niigata, NIIGATA	1.19	0.531	4.54	0.30	± 0.017	0.56	± 0.032	0.010	± 0.010
September, 1993									
Urawa, SAITAMA	1.68	0.389	7.96	0.12	± 0.016	0.30	± 0.040	0.057	± 0.0086

(5) Strontium-90 and Cesium-137 in Tea (Japanese Tea)

(from May. 1993 to Jun. 1993)

-continued from No. 103 of this publication-

Table (5) Strontium-90 and Cesium-137 in Tea (Japanese Tea)

Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kg)	(Bq/gCa)	(Bq/kg)	(Bq/kg)	(Bq/gK)	
May, 1993									
Nachikatsuura-machi, WAKAYAMA	5.10	2.36	20.0	2.3	± 0.09	0.96	± 0.040	0.69	± 0.053
Mifune-machi, KUMAMOTO	6.01	3.51	19.0	0.22	± 0.044	0.062	± 0.013	0.035	± 0.024
Kawaminami-machi, MIYAZAKI	4.79	2.95	17.8	0.90	± 0.079	0.30	± 0.027	1.3	± 0.08
Ikeda-machi, Gifu	4.60	2.15	17.2	0.77	± 0.052	0.36	± 0.024	0.18	± 0.036
Shirakawa-machi, Gifu	4.53	2.40	17.7	0.58	± 0.044	0.24	± 0.018	0.19	± 0.035
Iwata, SHIZUOKA	1.16	0.519	4.38	0.062	± 0.0089	0.12	± 0.017	0.046	± 0.0069
Shuzenji-machi, SHIZUOKA	1.55	1.04	5.63	1.6	± 0.04	1.6	± 0.04	0.20	± 0.013
Oodai-machi, MIE	5.06	2.30	19.1	0.46	± 0.040	0.20	± 0.017	0.32	± 0.043
Kameyama, MIE	5.07	2.38	18.6	0.58	± 0.045	0.24	± 0.019	0.12	± 0.032
Nara, NARA	4.81	2.36	18.5	0.51	± 0.041	0.21	± 0.018	0.13	± 0.030
Nara, NARA	5.92	2.68	22.4	0.26	± 0.029	0.096	± 0.011	0.12	± 0.028
Ue-mura, KUMAMOTO	4.79	3.63	15.7	0.80	± 0.074	0.22	± 0.020	0.31	± 0.045
Miyakonojou, MIYAZAKI	4.82	2.42	18.7	0.49	± 0.064	0.20	± 0.026	1.1	± 0.07
June, 1993									
Iruma, SAITAMA	5.36	2.56	19.1	0.55	± 0.044	0.22	± 0.017	0.40	± 0.050
Tokorozawa, SAITAMA	5.71	2.70	18.6	0.69	± 0.041	0.26	± 0.015	0.36	± 0.040
Uji, KYOTO	5.30	1.86	18.7	0.24	± 0.035	0.13	± 0.019	0.093	± 0.031
Kaya-machi, KYOTO	4.89	3.38	16.7	1.8	± 0.11	0.53	± 0.033	0.59	± 0.055
Chiran-machi, KAGOSHIMA	5.35	1.99	20.4	0.29	± 0.038	0.14	± 0.019	1.2	± 0.08
Miyanojou-machi, KAGOSHIMA	6.10	3.26	21.8	0.78	± 0.048	0.24	± 0.015	0.98	± 0.067

(6) Strontium-90 and cesium-137 in Sea Fish
(from Apr. 1993 to Sep. 1993)

-continued from No. 105 of this publication-

Table (6) :Strontium-90 and cesium-137 in Sea Fish

Location	Component			⁹⁰ Sr		¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)	(Bq/gCa)	(Bq/kgwet)	(Bq/gK)	
(Ammodytes personatus)								
April, 1993								
Akashi, HYOUGO (<i>Hexagrammos otakii</i>)	2.44	3.06	3.70	0.0008 ± 0.0035	0.0003 ± 0.0011	0.046 ± 0.0087	0.013 ± 0.0024	
September, 1993								
Fukushima, FUKUSHIMA (<i>Katsuwonus pelamis</i>)	1.79	2.35	4.21	0.0021 ± 0.0047	0.0009 ± 0.0020	0.18 ± 0.015	0.042 ± 0.0036	
May, 1993								
Tosa, KOCHI (<i>Limanda herzensteini</i>)	1.20	0.105	3.84	0.0039 ± 0.0062	0.037 ± 0.059	0.45 ± 0.025	0.12 ± 0.006	
June, 1993								
Sendai, MIYAGI (<i>Mugil cephalus</i>)	2.98	6.60	3.15	0.0028 ± 0.0062	0.00042 ± 0.00094	0.078 ± 0.0091	0.025 ± 0.0029	
September, 1993								
Morodomi-machi, SAGA (<i>Oncorhynchus keta</i>)	0.573	0.364	1.69	0.0016 ± 0.0071	0.004 ± 0.019	0.028 ± 0.0069	0.016 ± 0.0041	
September, 1993								
Urakawa-machi, HOKKAIDOU (<i>Pagrus sp</i>)	1.43	0.623	3.86	0.000 ± 0.010	0.000 ± 0.016	0.12 ± 0.011	0.031 ± 0.0028	
July, 1993								
Tennou-machi, AKITA	2.21	3.88	3.90	0.013 ± 0.0051	0.0033 ± 0.0013	0.16 ± 0.014	0.042 ± 0.0037	
Fukuoka, FUKUOKA (<i>Sardinops melanostictus</i>)	1.34	0.303	4.50	0.0000 ± 0.0056	0.000 ± 0.018	0.19 ± 0.013	0.043 ± 0.0028	
September, 1993								
Yamagata, YAMAGATA (<i>Scomber sp</i>)	2.68	6.26	2.49	0.0094 ± 0.0041	0.0015 ± 0.00065	0.069 ± 0.0096	0.028 ± 0.0038	

Location	Component			^{90}Sr		^{137}Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)	(Bq/gCa)	(Bq/kgwet)	(Bq/gK)	
August, 1993								
Matsuyama, EHIME <i>(Sebastiscus marmoratus)</i>	1.32	0.490	4.03	0.0057 ± 0.0067	0.012 ± 0.014	0.14 ± 0.011	0.034 ± 0.0028	
May, 1993								
Hamada, SHIMANE <i>(Sillago sp)</i>	5.98	18.4	2.55	0.027 ± 0.0054	0.0015 ± 0.00030	0.17 ± 0.015	0.066 ± 0.0057	
June, 1993								
Minamichita-machi, AICHI <i>(Trachurus sp)</i>	3.81	9.67	3.75	0.0000 ± 0.0058	0.00000 ± 0.00060	0.10 ± 0.010	0.027 ± 0.0028	
September, 1993								
Miyake-Island, TOKYO	1.47	1.61	3.53	0.0000 ± 0.0093	0.0000 ± 0.0058	0.18 ± 0.013	0.052 ± 0.0037	

Sea Fish

Japanese name	English name	Scientific name
Ikanago	Japanese sand lance	<u>Ammodytes personatus</u>
Ainame	Fat greenling	<u>Hexagrammos otakii</u>
Katsuo	Skipjack tuna	<u>Katsuwonus pelamis</u>
Magarei	Brown sole	<u>Limanda herzensteini</u>
Bora	Gray mullet	<u>Mugil cephalus</u>
Sake	Chum Salmon	<u>Oncorhynchus keta</u>
Tai	Sea bream	<u>Pagrus sp</u>
Maiwashi	Japanese pilchard	<u>Sardinops melanostictus</u>
Saba	Mackerel	<u>Scomber sp</u>
Kasago	Scorpion-fish	<u>Sebastiscus marmoratus</u>
Kisu	Whiting	<u>Sillago sp</u>
Aji	Horse mackerel	<u>Trachurus sp</u>

(7) Strontium-90 and cesium-137 in Freshwater Fish
 (from May. 1993 to Sep. 1993)

-continued from No. 105 of this publication-

Table (7) :Strontium-90 and cesium-137 in Freshwater Fish

Location	Component			⁹⁰ Sr				¹³⁷ Cs			
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)		(Bq/gCa)		(Bq/kgwet)		(Bq/gK)	
<u>(Carassius auratus)</u>											
July, 1993											
Ishikari-machi, HOKKAIDO	4.62	13.6	2.79	0.58	± 0.019	0.043	± 0.0014	0.086	± 0.011	0.031	± 0.0038
<u>(Cyprinus carpio)</u>											
May, 1993											
Kasumigaura-lake, IBARAKI	1.14	0.290	3.83	0.0087	± 0.0050	0.030	± 0.017	0.71	± 0.026	0.19	± 0.007
August, 1993											
Akita, AKITA	3.38	9.41	2.93	2.2	± 0.03	0.23	± 0.004	0.15	± 0.013	0.051	± 0.0043
September, 1993											
Fukushima, FUKUSHIMA	3.47	9.81	2.71	0.65	± 0.024	0.066	± 0.0024	0.20	± 0.018	0.074	± 0.0067

Freshwater Fish

Japanese name	English name	Scientific name
Fun	Crucian carp	<u>Carassius auratus</u>
Koi	Carp	<u>Cyprinus carpio</u>

(8) Strontium-90 and cesium-137 in Shellfish
 (from May, 1993 to Aug. 1993)

-continued from No. 105 of this publication-

Table (8) :Strontium-90 and cesium-137 in Shellfish

Location	Component			⁹⁰ Sr		¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)	(Bq/gCa)	(Bq/kgwet)	(Bq/gK)	
<u>(Mytilus edulis)</u>								
June, 1993								
Mutsu, AOMORI	2.73	0.624	1.52	0.0000 ± 0.0033	0.0000 ± 0.0052	0.023 ± 0.0064	0.015 ± 0.0042	
<u>(Ruditapes philippinarum)</u>								
May, 1993								
Konagai-machi, NAGASAKI	2.09	0.417	1.72	0.011 ± 0.0051	0.026 ± 0.012	0.0099 ± 0.0067	0.0057 ± 0.0039	
June, 1993								
Minamichita-machi, AICHI	2.13	0.893	3.52	0.0000 ± 0.0089	0.000 ± 0.010	0.060 ± 0.015	0.017 ± 0.0042	
<u>(Turbo cornutus)</u>								
May, 1993								
Ryotsu, NIIGATA	2.09	0.511	3.20	0.000 ± 0.010	0.000 ± 0.020	0.037 ± 0.016	0.012 ± 0.0049	
June, 1993								
Sakata, YAMAGATA	3.24	2.15	3.18	0.0025 ± 0.0039	0.0012 ± 0.0018	0.045 ± 0.0094	0.014 ± 0.0030	
August, 1993								
Togi-machi, ISHIKAWA	2.83	1.16	2.34	0.0000 ± 0.0042	0.0000 ± 0.0036	0.037 ± 0.0078	0.016 ± 0.0033	

Shellfish

Japanese name	English name	Scientific name
Murasakiigai	Common blue mussel	<u>Mytilus edulis</u>
Asari	Japanese littleneck	<u>Ruditapes philippinarum</u>
Sazae	Horned turban	<u>Turbo cornutus</u>

(9) Strontium-90 and cesium-137 in Seaweeds
 (from Apr. 1993 to May. 1993)

-continued from No. 105 of this publication-

Table (9) :Strontium-90 and cesium-137 in Seaweeds


Location	Component			⁹⁰ Sr				¹³⁷ Cs			
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)		(Bq/gCa)		(Bq/kgwet)		(Bq/gK)	
<u>(Undaria pinnatifida)</u>											
April, 1993											
Sakata, YAMAGATA	2.89	1.11	6.97	0.026	± 0.0051	0.023	± 0.0046	0.032	± 0.0077	0.0045	± 0.0011
Togi-machi, ISHIKAWA	3.83	0.878	5.78	0.013	± 0.0043	0.015	± 0.0049	0.040	± 0.0077	0.0069	± 0.0013
May, 1993											
Mutsu, AOMORI	3.04	0.793	6.75	0.019	± 0.0044	0.024	± 0.0056	0.021	± 0.0060	0.0032	± 0.00089
Fukaura-machi, AOMORI	2.87	0.814	6.56	0.030	± 0.0049	0.037	± 0.0060	0.028	± 0.0061	0.0042	± 0.00093
Ryotsu, NIIGATA	4.00	1.05	7.30	0.031	± 0.0064	0.029	± 0.0061	0.040	± 0.0084	0.0055	± 0.0011

Seaweeds

Japanese name	English name	Scientific name
Wakame	Wakame seaweed	<u>Undaria pinnatifida</u>

* * Total Diet * *

<Strontium-90>



<Cesium-137>





Fig. 1

* * Rice (producing districts) * *

<Strontium-90>



<Cesium-137>





Fig. 2

* * Milk (producing districts for domestic program)

<Strontium-90>



<Cesium-137>





Fig.3-1

* * Milk (producing districts for WHO program) * *

<Strontium-90>



<Cesium-137>





Fig. 3-2

* * Milk (consuming districts) * *

<Strontium-90>



<Cesium-137>





Fig. 3-3

* * Powdered Milk * *

<Strontium-90>



<Cesium-137>





Fig.3—4

* * Vegetables (producing districts) * *
(Japanese radish)

<Strontium-90>



<Cesium-137>





Fig. 4-1

* * Vegetables (consuming districts) * *
(Japanese radish)

<Strontium-90>



<Cesium-137>





Fig. 4-2

* * Tea (Japanese Tea) * *

<Strontium-90>



<Cesium-137>





Fig. 5

* * Sea Fish * *
(*Limanda herzensteini*)

<Strontium-90>



<Cesium-137>




Fig. 6





Fig. 7

* * Shellfish * *
(*Turbo cornutus*)

<Strontium-90>



<Cesium-137>

(Bq/kg wet)





Fig. 8

* * Seaweeds * *
(*Undaria pinnatifida*)

<Strontium-90>



<Cesium-137>




Fig. 9

* * Sampling Locations in Japan * *

- | | |
|----------------|---------------|
| 1: Sapporo | 36: Hiroshima |
| 2: Aomori | 37: Kochi |
| 3: Morioka | 38: Matsuyama |
| 4: Akita | 39: Yamaguchi |
| 5: Sendai | 40: Ooita |
| 6: Yamagata | 41: Fukuoka |
| 7: Fukushima | 42: Saga |
| 8: Niigata | 43: Kumamoto |
| 9: Mito | 44: Miyazaki |
| 10: Utsunomiya | 45: Nagasaki |
| 11: Chiba | 46: Kagoshima |
| 12: Urawa | 47: Naha |
| 13: Shinjuku | |
| 14: Maebashi | |
| 15: Nagano | |
| 16: Yokohama | |
| 17: Toyama | |
| 18: Kouhu | |
| 19: Kanazawa | |
| 20: Shizuoka | |
| 21: Gifu | |
| 22: Fukui | |
| 23: Nagoya | |
| 24: Tsu | |
| 25: Ootsu | |
| 26: Kyoto | |
| 27: Nara | |
| 28: Osaka | |
| 29: Tottori | |
| 30: Kobe | |
| 31: Wakayama | |
| 32: Okayama | |
| 33: Matsue | |
| 34: Tokushima | |
| 35: Takamatsu | |

