



ISSN 0441-2516

NIRS-RSD-117

RADIOACTIVITY SURVEY DATA in Japan

Part 2
= Dietary Materials =

NUMBER 117
September 1998

National Institute of Radiological Sciences
Chiba, Japan

Radioactivity Survey Data
in Japan
Number 117

September 1998 part 2 = Dietary Materials =

Contents

	Page
Environmental and Dietary Materials	
(Japan Chemical Analysis Center)	
1. Collection and pretreatment of samples	1
2. Preparation of samples for analysis	3
3. Separation of Strontium-90 and Cesium-137	3
4. Determination of Stable Strontium, Calcium and Potassium	4
5. Counting	4
6. Results	5
(1) Strontium-90 and Cesium-137 in Total Diet	5
(2) -1 Strontium-90 and Cesium-137 in Rice	9
(producing districts)	
-2 Strontium-90 and Cesium-137 in Rice	10
(consuming districts)	
(3) -1 Strontium-90 and Cesium-137 in Milk	12
(producing districts for domestic program)	
-2 Strontium-90 and Cesium-137 in Milk	13
(producing districts for WHO program)	
-3 Strontium-90 and Cesium-137 in Milk	14
(consuming districts)	
-4 Strontium-90 and Cesium-137 in Milk	16
(powdered milk)	
(4) -1 Strontium-90 and Cesium-137 in Vegetables	17
(producing districts)	
-2 Strontium-90 and Cesium-137 in Vegetables	21
(consuming districts)	
(5) Strontium-90 and Cesium-137 in Sea Fish	23
(6) Strontium-90 and Cesium-137 in Freshwater Fish	26
(7) Strontium-90 and Cesium-137 in Shellfish	28
(8) Strontium-90 and Cesium-137 in Seaweeds	30
7. Contents of Figure (Selected Location)	32

Environmental and Dietary Materials*

(Japan Chemical Analysis Center)

1. Collection and pretreatment of samples

(1) Rain and dry fallout

Rain and dry fallout was collected monthly on a sampling tray, approximately 5000cm² in area, which was filled with water to a depth of 1 cm at the beginning of every month.

Strontium and cesium carrier solutions were added after the sample was filtered. The tray was washed with 5 μ of distilled water and the washing was combined to the filtrate.

The sample was passed through a cation exchange column (500m μ of Dowex 50W X8, 50~100 mesh, Na form) at a rate flow of 80m μ /min.

(2) Airborne dust

Airborne dust was collected by an electrostatic precipitator or a filter air sampler for every three-months at a rate of more than 3000m³ per month.

The sampling was done 1 to 1.5 meters above the ground.

(3) Service water and freshwater

Service water, 100 μ each, was collected at the intake of the water-treatment plant and at the tap after water was left running for five minutes. Strontium and cesium carriers were added to the filtered water sample. The subsequent process was the same as that described in the section (1). Freshwater was treated in the same way as the service water.

(4) Soil

Soil was collected from the location in the spacious and flat area without past surface disturbance caused by dust storms, inflow and out flow due to precipitation, etc.. Any places located under trees in a forest, in a stony area or inside of river banks were avoided. Soil was taken from two layers of different depths, 0-5cm and 5-20cm. The soil lumps were crushed by hands and dried in a drying oven regulated 105°C . The soil was then passed through a 2mm sieve to remove plant roots and pebbles.

(5) Sea water

Sea water was collected at the fixed stations

where the effect of terrestrial fresh water from rivers was expected to be negligibly small. A special consideration was also given to weather conditions.

The sampling was carried out when there was no rainfall for the last few days. To prevent contamination, water samples were collected at the bow of a sampling boat just before she stood still by scooping surface water using a polyethylene bucket.

Immediately after the collection, the samples were acidified to a pH lower than 3 by adding concentrated hydrochloric acid in a ratio of 1m μ to 1 μ of sea water, and then stored in 20 μ polyethylene containers. The sampling equipments as well as containers were thoroughly rinsed with dilute hydrochloric acid and then with distilled water before use. Two hundred milliliters of sea water was also collected at the same stations for the determination of chlorinity.

(6) Sea sediments

Sediment was collected in the same area as that for the sea water sample, taking the following criteria into account:

- a. The depth of water exceeds 1m at low tide.
- b. No significant sedimental movement is observed in the vicinity of concern.
- c. Mud, silt and fine sand are preferable.

A conventional sediment sampling device was used for collecting the top few centimeters of surface sediment. Approximately 4kg of the sample in wet weight was spread on a stainless steel dish after removed of the pebbles, shells and other foreign materials, and dried in a drying oven regulated at 105°C.

(7) Total diet

A full one day ordinary diet including three meals, water, tea and other in-between snacks for five persons was collected as a sample of "total diet".

The sample in a large stainless steel pan was carbonized carefully by direct application of gas flame, and was transferred to a porcelain dish and then ashed at 450°C in an electric muffle furnace.

(8) Rice

Polished rice was collected in producing districts at the harvest and in consuming areas when new crops were first put on sale. The sample was carbonized and ashed in a porcelain dish.

* Samples were sent to the Center from 46 contracted prefectures.

(9) Milk

Raw milk was collected in producing districts and commercial milk was purchased in consuming districts. Milk in a stainless steel pan or a porcelain dish was evaporated to dryness followed by carbonization and ashing.

(10) Vegetables

Spinach and Japanese radish were selected as the representatives for leaf vegetables and for nonstarch roots, respectively. After removing soil, the edible part of vegetable sample was dried and carbonized in a stainless steel pan or a porcelain dish.

(11) Tea

Five hundred grams of manufactured green tea was collected, carbonized and ashed in a stainless steel pan or a porcelain dish.

(12) Fish, shellfish and seaweeds

a. Sea fish and freshwater fish

Fish was rinsed with water and blotted with a filter paper. Only the edible part was used in case of larger sized fish, and the whole part was used in case of smaller ones. Each sample was weighed and placed in a stainless steel pan or a porcelain dish. After carbonized, the sample was ashed in an electric muffle furnace.

b. Shellfish

Approximately 4kg of shellfish including the shells was collected or purchased. After removing the shells, it was treated in the same way as that for the sea fish.

c. Seaweeds

Edible seaweeds were collected and rinsed with water to remove sand and other adhering matters on the surface. These were removed of excess water, weighed dried and ashed.

Table 1 shows details of sample collection.

Table 1 Details of sample collection

Sample	Frequency of sampling	Quantity of sample
=Environmental materials=		
(1) Rain and dry fallout		
1. For domestic program	monthly	
2. For WHO program	monthly	
(2) Airborne dust	quarterly	>3000 m ³ /month
(3) Service water and freshwater		
1. Service water (source water)	semiyearly	100 ℥
2. Service water (tap water)	semiyearly	100 ℥
3. Freshwater	yearly (fishing season)	100 ℥
(4) Soil		
1. 0~ 5 cm	yearly	4 kg
2. 5~ 20cm	yearly	4 kg
(5) Sea water	yearly	40 ℥
(6) Sea sediments	yearly	4 kg
=Dietary materials=		
(7) Total diet	semiyearly	daily amount for 5 persons
(8) Rice		
1. Producing districts	yearly (harvesting season)	5 kg (polished rice)
2. Consuming districts	yearly (harvesting season)	5 kg (polished rice)
(9) Milk		
1. Producing districts for WHO program	quarterly (February, May, August and November)	3 ℥
2. Producing districts for domestic program	semiyearly (February and August)	3 ℥

Sample	Frequency of sampling	Quantity of sample
3. Consuming districts	semiyearly (February and August)	3 ℥
4. Powdered milk	semiyearly (April and October)	2~ 3 kg
(10) Vegetables		
1. Producing districts	yearly (harvesting season)	4 kg
2. Consuming districts	yearly (harvesting season)	4 kg
(11) Tea	yearly (the first harvesting season)	500g (manufactured tea)
(12) Fish, shellfish and seaweeds		
1. Sea fish	yearly (fishing season)	4 kg
2. Freshwater fish	yearly (fishing season)	4 kg
3. Shellfish	yearly (fishing season)	4 kg
4. Seaweeds	yearly (fishing season)	2~ 3 kg

2. Preparation of samples for analysis

(1) Rain, service water and freshwater

Strontium and cesium were eluted with hydrochloric acid from the cation exchange column. The residue of rain sample on the filter paper was ashed in an electric muffle furnace and the ash was dissolved in hydrochloric acid. The insoluble part was filtered and washed. The filtrate and the washings were combined to the previous eluate and used for radiochemical analysis.

(2) Soil and Sea sediment

Dried soil was crushed to smaller ones than 0.2 5mm in size by a crusher. The sieved sample was ashed in an electric muffle furnace regulated at 450 °C. The sample was then heated with hydrochloric acid, strontium and cesium carrier solutions and the mixture was heated. The insoluble constituent was filtered off and washed with water.

The dried sample was crushed to smaller ones than 0.25mm by a crushing machine. The further preparation of the sample was the same as that described in the section 2-(2).

(3) Rice

The ashed sample was pulverized with a porcelain mortar and passed through a 0.35mm sieve. The sieved sample to which both strontium and cesium carriers were added, was digested with nitric acid by heating. After the sample was heated again with nitric acid to dryness, strontium and cesium were extracted with hydrochloric acid and water. The insoluble constituent was filtered and washed. The filtrate and washings were combined for subsequent radiochemical analysis.

(4) Airborne dust, diet, milk, vegetables, fish and shellfish, seaweeds, tea and others

These ashed samples were treated with the

same procedure as that described in the section 2-(4).

3. Separation of strontium-90 and cesium-137

(1) Strontium-90

Sample solutions, prepared as in the foregoing sections 2-(1) through 2-(4), were neutralized with sodium hydroxide. After sodium carbonate was added, the precipitate of strontium and calcium carbonates was separated. The supernatant solution was retained for cesium-137 determination.

The carbonates were dissolved in hydrochloric acid and strontium and calcium were precipitated as oxalates. The precipitate was dissolved in nitric acid and strontium was separated from calcium by successive fuming nitric acid separation. Iron scavenge was made after addition of ferric iron carrier followed by barium chromate separation after addition of barium carrier to remove radium, its daughters and lead. Strontium was recovered as carbonate, and the precipitate was dried and weighed to determine strontium recovery. The strontium carbonate was dissolved in hydrochloric acid and iron carrier was added. The solution was allowed to stand for two weeks for strontium-90 and yttrium-90 to attain equilibrium. Yttrium-90 was coprecipitated with ferric hydroxide and the precipitate was filtered off, washed and counted.

(2) Cesium-137

The supernatant separated from the strontium fraction was acidified with hydrochloric acid. While stirring, cesium was adsorbed on the ammonium molybdate phosphate added.

After filtered off and washed with hydrochloric acid the precipitate was dissolved in 2.5N sodium hydroxide solution. The solution was adjusted to pH 8.2 with hydrochloric acid and allowed to cool.

Resultant molybdenum hydroxide which separated

out in the solution, was filtered off and washed with water. EDTA was added to the filtrate and washings. Cesium and rubidium were adsorbed on a cation exchange column and cesium was separated from rubidium by eluting with hydrochloric acid.

The eluate was evaporated to dryness and was dissolved. The solution was filtered.

Chloroplatinic acid was added to precipitate cesium. The precipitate was filtered onto a tared paper using a demountable filter and washed with water and then ethanol. After drying, the chemical yield of cesium was determined by weighing the precipitate. Cesium-137 radioactivity was measured for this precipitate.

4. Determination of stable strontium, calcium and potassium

A weighed amount of soil or sea sediment was heated in a electric muffle furnace at 450°C and then treated with hydrochloric acid for extraction. A weighed aliquot of ashed samples of total diet, vegetables, milk, fish, shellfish or seaweeds was

digested with hydrofluoric acid and nitric acid. The extract was made up to an appropriate volume with dilute hydrochloric acid. The sample solution was analyzed for calcium by titration with standard potassium permanganate solution after separating calcium as oxalate. Atomic absorption spectroscopy was applied when appropriate. Stable strontium and potassium were determined by atomic absorption and flame emission spectrometry, respectively.

5. Counting

After the radiochemical separation the mounted precipitates were counted for activity using low background beta counters normally for 60 to 90min. Net sample counting rates were corrected for counter efficiency, recovery, self-absorption and decay to obtain the content of strontium-90 and cesium-137 per sample aliquot. From the results, concentrations of these nuclides in the original samples were calculated.

6. Results

(1) Strontium-90 and Cesium-137 in Total Diet

(from Oct. 1995 to Mar. 1996)

-continued from No. 115 of this publication-

Table (1) Strontium-90 and Cesium-137 in Total Diet

Location	Ash	Ca	K	⁹⁰ Sr				¹³⁷ Cs			
	Ash(g/p·d)	Ca(mg/p·d)	K(mg/p·d)	(Bq/p·d)	(Bq/gCa)	(Bq/p·d)	(Bq/gK)				
October, 1995											
Kamiita-machi, TOKUSHIMA	18.6	623	2310	0.050	± 0.0055	0.080	± 0.0088	0.029	± 0.0075	0.013	± 0.0032
Kochi, KOCHI	13.4	488	1890	0.052	± 0.0061	0.11	± 0.012	0.020	± 0.0057	0.011	± 0.0030
Saga-machi, KOCHI	15.2	464	2020	0.069	± 0.0070	0.15	± 0.015	0.021	± 0.0075	0.010	± 0.0037
Ooita, OITA	11.8	406	1840	0.052	± 0.0060	0.13	± 0.015	0.052	± 0.0075	0.028	± 0.0041
Saiki, OITA	13.7	498	1560	0.046	± 0.0061	0.093	± 0.012	0.015	± 0.0054	0.0099	± 0.0035
November, 1995											
Iwajumi-machi, IWATE	13.2	344	1720	0.058	± 0.0070	0.17	± 0.020	0.063	± 0.0090	0.037	± 0.0053
Ishinomaki, MIYAGI	17.4	533	2200	0.076	± 0.0070	0.14	± 0.013	0.043	± 0.0082	0.020	± 0.0037
Onagawa-machi, MIYAGI	20.0	608	2330	0.059	± 0.0065	0.098	± 0.011	0.040	± 0.0089	0.017	± 0.0038
Yamagata, YAMAGATA	16.6	500	1710	0.041	± 0.0061	0.081	± 0.012	0.064	± 0.0083	0.038	± 0.0049
Sagae, YAMAGATA	13.8	376	1540	0.047	± 0.0056	0.12	± 0.015	0.029	± 0.0073	0.019	± 0.0048
Fukushima, FUKUSHIMA	14.3	337	1790	0.050	± 0.0055	0.15	± 0.016	0.020	± 0.0047	0.011	± 0.0026
Ookuma-machi, FUKUSHIMA	11.7	716	1430	0.060	± 0.0068	0.084	± 0.0095	0.036	± 0.0065	0.025	± 0.0046
Utsunomiya, TOCHIGI	14.4	505	2180	0.069	± 0.0065	0.14	± 0.013	0.062	± 0.0086	0.029	± 0.0039
Mooka, TOCHIGI	12.8	408	1590	0.056	± 0.0062	0.14	± 0.015	0.039	± 0.0075	0.025	± 0.0047
Yokohama, KANAGAWA	16.5	485	2300	0.046	± 0.0061	0.095	± 0.013	0.053	± 0.0087	0.023	± 0.0038
Hiratsuka, KANAGAWA	15.1	606	2200	0.052	± 0.0064	0.086	± 0.011	0.049	± 0.0084	0.022	± 0.0038
Takaoka, TOYAMA	12.0	266	1600	0.047	± 0.0089	0.17	± 0.034	0.042	± 0.0063	0.026	± 0.0040
Takaoka, TOYAMA	11.5	365	1720	0.056	± 0.0061	0.15	± 0.017	0.080	± 0.0085	0.047	± 0.0050
Fukui, FUKUI	15.2	550	1830	0.063	± 0.0067	0.11	± 0.012	0.026	± 0.0073	0.014	± 0.0040
Shizuoka, SHIZUOKA	16.1	695	2590	0.064	± 0.011	0.092	± 0.015	0.054	± 0.0074	0.021	± 0.0029
Hamaoka-machi, SHIZUOKA	14.2	571	1930	0.045	± 0.0098	0.078	± 0.017	0.017	± 0.0047	0.0089	± 0.0024
Nagoya, AICHI	18.7	642	2530	0.087	± 0.012	0.14	± 0.019	0.053	± 0.0087	0.021	± 0.0034

Location	Ash	Ca	K	^{90}Sr				^{137}Cs		
	Ash(g/p·d)	Ca(mg/p·d)	K(mg/p·d)	(Bq/p·d)	(Bq/gCa)	(Bq/p·d)	(Bq/gK)			
Shinshiro, AICHI	12.0	580	1590	0.024 ± 0.0078	0.041 ± 0.013	0.035 ± 0.0072	0.022 ± 0.0045			
Tsu, MIE	20.0	779	2210	0.072 ± 0.0069	0.092 ± 0.0088	0.042 ± 0.0087	0.019 ± 0.0039			
Kashihara, NARA	13.5	785	1700	0.076 ± 0.0077	0.097 ± 0.0097	0.075 ± 0.0095	0.044 ± 0.0056			
Gojou, NARA	13.1	1450	1470	0.061 ± 0.010	0.042 ± 0.0069	0.0096 ± 0.0059	0.0065 ± 0.0040			
Wakayama, WAKAYAMA	13.4	468	1780	0.055 ± 0.0062	0.12 ± 0.013	0.037 ± 0.0073	0.021 ± 0.0041			
Kitayama-mura, WAKAYAMA	15.3	577	1680	0.076 ± 0.0071	0.13 ± 0.012	0.035 ± 0.0067	0.021 ± 0.0040			
Kashima-machi, SHIMANE	15.9	642	2240	0.068 ± 0.0070	0.11 ± 0.011	0.041 ± 0.0080	0.018 ± 0.0035			
Okayama, OKAYAMA	18.5	654	2310	0.072 ± 0.0070	0.11 ± 0.011	0.063 ± 0.0080	0.027 ± 0.0034			
Kamisaibara-mura, OKAYAMA	16.4	419	2070	0.10 ± 0.009	0.25 ± 0.021	0.086 ± 0.0091	0.041 ± 0.0044			
Matsuyama, EHIME	12.2	536	1760	0.048 ± 0.0059	0.090 ± 0.011	0.030 ± 0.0060	0.017 ± 0.0034			
Ikata-machi, EHIME	9.49	413	1230	0.041 ± 0.0053	0.099 ± 0.013	0.032 ± 0.0056	0.026 ± 0.0045			
Dazaifu, FUKUOKA	12.6	549	1930	0.053 ± 0.0062	0.097 ± 0.011	0.044 ± 0.0082	0.023 ± 0.0042			
Saga, SAGA	17.0	751	2400	0.080 ± 0.0073	0.11 ± 0.010	0.044 ± 0.0077	0.018 ± 0.0032			
Nagasaki, NAGASAKI	15.7	479	2050	0.062 ± 0.0064	0.13 ± 0.013	0.049 ± 0.0070	0.024 ± 0.0034			
Matsuura, NAGASAKI	13.3	464	1640	0.029 ± 0.0050	0.063 ± 0.011	0.046 ± 0.0073	0.028 ± 0.0044			
Sendai, MIYAGI	12.5	361	1330	0.036 ± 0.0057	0.098 ± 0.016	0.039 ± 0.0077	0.029 ± 0.0058			
Ookuchi, KAGOSHIMA	17.4	517	2020	0.067 ± 0.0072	0.13 ± 0.014	0.045 ± 0.0077	0.022 ± 0.0038			
December, 1995										
Sapporo, HOKKAIDOU	14.6	711	2510	0.081 ± 0.0074	0.11 ± 0.010	0.023 ± 0.0075	0.0092 ± 0.0030			
Iwanai-machi, HOKKAIDOU	12.9	454	1780	0.046 ± 0.0060	0.10 ± 0.013	0.050 ± 0.0088	0.028 ± 0.0050			
Aomori, AOMORI	16.5	417	2090	0.057 ± 0.0063	0.14 ± 0.015	0.079 ± 0.0087	0.038 ± 0.0042			
Ajigasawa-machi, AOMORI	15.3	544	2560	0.074 ± 0.0072	0.14 ± 0.013	0.036 ± 0.0066	0.014 ± 0.0026			
Morioka, IWATE	16.0	577	1810	0.060 ± 0.0069	0.10 ± 0.012	0.042 ± 0.0078	0.023 ± 0.0043			
Akita, AKITA	18.1	430	2090	0.077 ± 0.0075	0.18 ± 0.018	0.10 ± 0.010	0.048 ± 0.0050			
Akita, AKITA	14.0	440	1800	0.061 ± 0.0071	0.14 ± 0.016	0.060 ± 0.0088	0.033 ± 0.0049			

Location	Ash	Ca	K	^{90}Sr				^{137}Cs		
	Ash(g/p·d)	Ca(mg/p·d)	K(mg/p·d)	(Bq/p·d)	(Bq/gCa)	(Bq/p·d)	(Bq/gK)			
Mito, IBARAKI	16.1	796	2420	0.044	\pm 0.0091	0.055	\pm 0.011	0.044	\pm 0.0067	0.018 \pm 0.0028
Tokai-mura, IBARAKI	20.5	737	3250	0.10	\pm 0.012	0.14	\pm 0.017	0.095	\pm 0.010	0.029 \pm 0.0031
Maebashi, GUNMA	13.3	413	2070	0.059	\pm 0.0059	0.14	\pm 0.014	0.062	\pm 0.0091	0.030 \pm 0.0044
Nakanojou-machi, GUNMA	18.5	946	2520	0.076	\pm 0.0069	0.080	\pm 0.0073	0.048	\pm 0.0086	0.019 \pm 0.0034
Urawa, SAITAMA	16.6	950	2210	0.073	\pm 0.0070	0.077	\pm 0.0074	0.079	\pm 0.0098	0.036 \pm 0.0044
Kumagaya, SAITAMA	15.3	645	2010	0.052	\pm 0.0062	0.080	\pm 0.0097	0.046	\pm 0.0081	0.023 \pm 0.0040
Ichihara, CHIBA	15.2	451	1660	0.033	\pm 0.0093	0.073	\pm 0.021	0.056	\pm 0.010	0.034 \pm 0.0060
Chikura-machi, CHIBA	19.9	693	2450	0.11	\pm 0.024	0.16	\pm 0.035	0.007	\pm 0.016	0.0029 \pm 0.0065
Shinjuku, TOKYO	13.9	440	1610	0.041	\pm 0.0054	0.092	\pm 0.012	0.024	\pm 0.0078	0.015 \pm 0.0048
Hachijou-machi, TOKYO	16.0	934	2210	0.057	\pm 0.0059	0.061	\pm 0.0063	0.070	\pm 0.010	0.032 \pm 0.0047
Nishikawa-machi, NIIGATA	23.4	718	3110	0.078	\pm 0.011	0.11	\pm 0.015	0.034	\pm 0.0062	0.011 \pm 0.0020
Kashiwazaki, NIIGATA	21.0	557	2330	0.073	\pm 0.011	0.13	\pm 0.019	0.031	\pm 0.0063	0.013 \pm 0.0027
Kanazawa, ISHIKAWA	15.7	628	2130	0.055	\pm 0.0062	0.087	\pm 0.0099	0.049	\pm 0.0086	0.023 \pm 0.0040
Yoshinodani-mura, ISHIKAWA	15.2	418	1620	0.040	\pm 0.0056	0.095	\pm 0.014	0.049	\pm 0.0088	0.030 \pm 0.0054
Koufu, YAMANASHI	14.9	337	2000	0.059	\pm 0.0069	0.17	\pm 0.021	0.025	\pm 0.0054	0.013 \pm 0.0027
Nirasaki, YAMANASHI	16.5	706	2200	0.056	\pm 0.0097	0.080	\pm 0.014	0.035	\pm 0.0062	0.016 \pm 0.0028
Nagano, NAGANO	17.4	736	2520	0.055	\pm 0.0065	0.075	\pm 0.0088	0.016	\pm 0.0065	0.0063 \pm 0.0026
Sanada-machi, NAGANO	15.9	642	2110	0.059	\pm 0.0064	0.092	\pm 0.010	0.030	\pm 0.0070	0.014 \pm 0.0033
Gifu, GIFU	13.9	528	2190	0.052	\pm 0.0065	0.099	\pm 0.012	0.046	\pm 0.0082	0.021 \pm 0.0038
Takayama, GIFU	13.8	502	2040	0.051	\pm 0.0066	0.10	\pm 0.013	0.031	\pm 0.0072	0.015 \pm 0.0035
Ootsu, SHIGA	15.8	607	2070	0.055	\pm 0.0062	0.091	\pm 0.010	0.038	\pm 0.0076	0.018 \pm 0.0037
Imazu-machi, SHIGA	19.1	1000	2550	0.16	\pm 0.010	0.16	\pm 0.010	0.083	\pm 0.010	0.033 \pm 0.0040
Owase, MIE	13.9	465	1660	0.055	\pm 0.0061	0.12	\pm 0.013	0.028	\pm 0.0077	0.017 \pm 0.0047
Kyoto, KYOTO	17.8	528	2450	0.045	\pm 0.0055	0.086	\pm 0.010	0.047	\pm 0.0072	0.019 \pm 0.0029
Maizuru, KYOTO	18.4	599	2440	0.057	\pm 0.0062	0.096	\pm 0.010	0.042	\pm 0.0066	0.017 \pm 0.0027

Location	Ash	Ca	K	⁹⁰ Sr				¹³⁷ Cs		
	Ash(g/p·d)	Ca(mg/p·d)	K(mg/p·d)	(Bq/p·d)	(Bq/gCa)	(Bq/p·d)	(Bq/gK)			
Osaka, OSAKA	13.3	487	2000	0.031 ± 0.0080	0.065 ± 0.016	0.035 ± 0.0073	0.018 ± 0.0036			
Sakai, OSAKA	14.8	522	2360	0.062 ± 0.0067	0.12 ± 0.013	0.053 ± 0.0075	0.022 ± 0.0032			
Kakogawa, HYOGO	12.1	598	1600	0.045 ± 0.0058	0.074 ± 0.0098	0.017 ± 0.0070	0.011 ± 0.0044			
Hamasaka-machi, HYOGO	13.6	576	1790	0.043 ± 0.0056	0.074 ± 0.0098	0.027 ± 0.0072	0.015 ± 0.0041			
Matsue, SHIMANE	23.0	1180	3340	0.12 ± 0.009	0.10 ± 0.007	0.053 ± 0.0090	0.016 ± 0.0027			
Miyoshi, HIROSHIMA	12.0	426	1410	0.030 ± 0.0085	0.070 ± 0.020	0.035 ± 0.0063	0.025 ± 0.0045			
Yamaguchi, YAMAGUCHI	15.5	549	2230	0.070 ± 0.011	0.13 ± 0.020	0.048 ± 0.0080	0.022 ± 0.0036			
Ajisui-machi, YAMAGUCHI	18.0	597	2250	0.044 ± 0.0055	0.075 ± 0.0092	0.060 ± 0.0082	0.027 ± 0.0037			
Tokushima, TOKUSHIMA	17.7	544	2630	0.049 ± 0.0054	0.090 ± 0.010	0.035 ± 0.0077	0.013 ± 0.0029			
Takamatsu, KAGAWA	14.5	401	2030	0.041 ± 0.0097	0.10 ± 0.024	0.015 ± 0.0047	0.0076 ± 0.0023			
Tuda-machi, KAGAWA	13.8	482	2100	0.070 ± 0.011	0.15 ± 0.023	0.030 ± 0.0064	0.014 ± 0.0030			
Fukuoka, FUKUOKA	14.2	291	1030	0.040 ± 0.0054	0.14 ± 0.019	0.049 ± 0.0082	0.048 ± 0.0079			
Karatsu, SAGA	16.1	476	1960	0.050 ± 0.0062	0.10 ± 0.013	0.031 ± 0.0071	0.016 ± 0.0036			
Kumamoto, KUMAMOTO	13.3	351	2000	0.052 ± 0.0061	0.15 ± 0.017	0.038 ± 0.0068	0.019 ± 0.0034			
Aso-machi, KUMAMOTO	14.3	376	1890	0.070 ± 0.0068	0.19 ± 0.018	0.048 ± 0.0073	0.026 ± 0.0039			
Miyazaki, MIYAZAKI	17.9	567	2590	0.066 ± 0.0067	0.12 ± 0.012	0.041 ± 0.0070	0.016 ± 0.0027			
Takahara-machi, MIYAZAKI	12.6	262	1600	0.045 ± 0.0055	0.17 ± 0.021	0.078 ± 0.0085	0.048 ± 0.0053			
Naha, Okinawa	15.7	512	2510	0.052 ± 0.0063	0.10 ± 0.012	0.035 ± 0.0069	0.014 ± 0.0027			
January, 1996										
Tsuruga, FUKUI	15.4	586	1740	0.068 ± 0.0064	0.12 ± 0.011	0.032 ± 0.0071	0.018 ± 0.0041			
Hirosima, HIROSHIMA	12.7	408	1700	0.057 ± 0.010	0.14 ± 0.025	0.021 ± 0.0050	0.012 ± 0.0029			
Febraly, 1996										
Kamiita-machi, TOKUSHIMA	17.9	482	2530	0.039 ± 0.0084	0.082 ± 0.018	0.031 ± 0.0060	0.012 ± 0.0024			
Ginowan, Okinawa	16.0	633	2230	0.053 ± 0.0064	0.083 ± 0.010	0.047 ± 0.0084	0.021 ± 0.0038			

(2)-1 Strontium-90 and Cesium-137 in Rice (producing districts)

(from Oct. 1995 to Mar. 1996)

-continued from No. 115 of this publication-

Table (2)-1 Strontium-90 and Cesium-137 in Rice

Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kg wet)	(Bq/g Ca)		(Bq/kg wet)	(Bq/g K)	
October, 1995									
Maki-machi, NIIGATA	0.529	0.021	0.746	0.0031 ± 0.0064	0.15 ± 0.30		0.0014 ± 0.0040	0.0019 ± 0.0054	
Kosugi-machi, TOYAMA	0.505	0.028	0.853	0.0007 ± 0.0062	0.02 ± 0.23		0.017 ± 0.0053	0.020 ± 0.0062	
Shiga-machi, SHIGA	0.568	0.043	0.829	0.0048 ± 0.0032	0.11 ± 0.075		0.026 ± 0.0070	0.032 ± 0.0085	
Kashihara, NARA	0.723	0.035	1.09	0.0043 ± 0.0028	0.12 ± 0.079		0.012 ± 0.0055	0.011 ± 0.0051	
Ishii-machi, TOKUSHIMA	0.411	0.025	0.801	0.0000 ± 0.0047	0.00 ± 0.19		0.0000 ± 0.0028	0.0000 ± 0.0034	
Miki-machi, KAGAWA	0.520	0.036	0.738	0.0012 ± 0.0030	0.035 ± 0.084		0.0000 ± 0.0032	0.0000 ± 0.0043	
Koushi-machi, KUMAMOTO	0.507	0.024	0.649	0.0034 ± 0.0031	0.14 ± 0.13		0.0000 ± 0.0046	0.0000 ± 0.0070	
Usa, OITA	0.637	0.029	1.11	0.0055 ± 0.0038	0.19 ± 0.13		0.0000 ± 0.0036	0.0000 ± 0.0032	
November, 1995									
Ishikari-machi, HOKKAIDOU	0.655	0.039	0.917	0.0018 ± 0.0033	0.046 ± 0.084		0.0069 ± 0.0043	0.0075 ± 0.0047	
Takizawa-mura, IWATE	0.644	0.032	1.04	0.0007 ± 0.0028	0.023 ± 0.087		0.21 ± 0.014	0.20 ± 0.013	
Ishinomaki, MIYAGI	0.552	0.040	0.745	0.0085 ± 0.0032	0.21 ± 0.079		0.0000 ± 0.0052	0.0000 ± 0.0070	
Fukushima, FUKUSHIMA	0.753	0.040	1.20	0.0046 ± 0.0034	0.12 ± 0.087		0.0080 ± 0.0046	0.0066 ± 0.0038	
Mito, IBARAKI	0.475	0.044	0.627	0.0003 ± 0.0032	0.006 ± 0.074		0.036 ± 0.0064	0.057 ± 0.010	
Shinguu, WAKAYAMA	0.591	0.025	1.32	0.0000 ± 0.0036	0.00 ± 0.14		0.0028 ± 0.0038	0.0021 ± 0.0029	
Saga, SAGA	0.588	0.033	0.694	0.0005 ± 0.0031	0.016 ± 0.094		0.0079 ± 0.0060	0.011 ± 0.0086	
December, 1995									
Utsunomiya, TOCHIGI	0.655	0.034	0.779	0.0090 ± 0.0037	0.26 ± 0.11		0.010 ± 0.0059	0.013 ± 0.0076	
Maebashi, GUNMA	0.657	0.040	0.788	0.0058 ± 0.0030	0.15 ± 0.076		0.011 ± 0.0058	0.014 ± 0.0074	
Takane-machi, YAMANASHI	0.842	0.023	1.27	0.0022 ± 0.0043	0.1 ± 0.19		0.0074 ± 0.0044	0.0058 ± 0.0035	
Kasai, HYOUNGO	0.576	0.032	0.657	0.014 ± 0.0037	0.43 ± 0.12		0.0000 ± 0.0050	0.0000 ± 0.0076	
Yamaguchi, YAMAGUCHI	0.735	0.041	1.17	0.0078 ± 0.0035	0.19 ± 0.086		0.075 ± 0.0091	0.064 ± 0.0078	
January, 1996									
Chikushino, FUKUOKA	0.654	0.027	0.929	0.0016 ± 0.0037	0.06 ± 0.14		0.075 ± 0.0090	0.081 ± 0.0097	

(10)

(2)-2 Strontium-90 and Cesium-137 in Rice (consuming districts)
(from Oct. 1995 to Mar. 1996)
-continued from No. 113 of this publication-
Table (2)-2 Strontium-90 and Cesium-137 in Rice

Location	Component			⁸⁹ Sr			¹³⁷ Cs	
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)	(Bq/gCa)	(Bq/kgwet)	(Bq/gK)	
Hirosaki, AOMORI	0.586	0.032	1.04	0.0045 ± 0.0063	0.14 ± 0.20	0.0088 ± 0.0044	0.0085 ± 0.0042	
Nagasaki, NAGASAKI	0.528	0.023	0.908	0.0021 ± 0.0065	0.09 ± 0.28	0.015 ± 0.0051	0.016 ± 0.0056	

(3)-1 Strontium-90 and Cesium-137 in Milk (producing districts for domestic program)

(from Oct. 1995 to Mar. 1996)

-continued from No. 115 of this publication-

Table (3)-1 Strontium-90 and Cesium-137 in Milk

Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(g/l)	Ca(g/l)	K(g/l)	(Bq/l)	(Bq/gCa)		(Bq/l)	(Bq/gK)	
October, 1995									
Yamato-machi, SAGA	7.22	1.09	1.63	0.029	± 0.0050	0.027	± 0.0046	0.036	± 0.0064
January, 1996									
Shinguu, WAKAYAMA	6.80	1.02	1.50	0.013	± 0.0061	0.013	± 0.0060	0.013	± 0.0057
February, 1996									
Aomori, AOMORI	7.48	1.10	1.57	0.048	± 0.0066	0.043	± 0.0061	0.030	± 0.0066
Takizawa-mura, IWATE	7.08	1.04	1.54	0.021	± 0.0044	0.020	± 0.0043	0.060	± 0.0082
Mito, IBARAKI	7.41	1.15	1.58	0.079	± 0.0066	0.069	± 0.0058	0.016	± 0.0058
Nishinasuno-machi, TOCHIGI	7.81	1.21	1.74	0.026	± 0.0047	0.021	± 0.0039	0.039	± 0.0081
Fujimi-mura, GUNMA	6.68	1.02	1.39	0.023	± 0.0046	0.023	± 0.0045	0.019	± 0.0053
Yachimata, CHIBA	7.91	1.19	1.63	0.029	± 0.0054	0.025	± 0.0045	0.021	± 0.0059
Tonami, TOYAMA	7.37	1.09	1.51	0.017	± 0.0051	0.016	± 0.0047	0.067	± 0.0086
Oshimizu-machi, ISHIKAWA	7.20	1.11	1.57	0.030	± 0.0059	0.027	± 0.0053	0.030	± 0.0068
Kasamatsu-machi, GIFU	6.71	1.02	1.38	0.040	± 0.0059	0.039	± 0.0058	0.0023	± 0.0041
Hino-machi, SHIGA	7.21	1.09	1.52	0.012	± 0.0061	0.011	± 0.0056	0.0048	± 0.0046
Oouchiyama-mura, MIE	7.42	1.11	1.59	0.012	± 0.0063	0.011	± 0.0057	0.017	± 0.0059
Mihara-machi, HYOUGO	6.60	1.000	1.44	0.023	± 0.0074	0.023	± 0.0074	0.0082	± 0.0057
Oouda-machi, NARA	7.25	1.03	1.58	0.028	± 0.0087	0.028	± 0.0084	0.0034	± 0.0046
Takase-machi, KAGAWA	7.19	1.10	1.55	0.029	± 0.0051	0.027	± 0.0046	0.0063	± 0.0047
Matsuyama, EHIME	7.34	1.11	1.53	0.026	± 0.0051	0.024	± 0.0046	0.011	± 0.0051
Kujuu-machi, OOITA	7.13	1.11	1.55	0.019	± 0.0045	0.017	± 0.0041	0.11	± 0.011
Takahara-machi, MIYAZAKI	7.35	1.08	1.61	0.041	± 0.0091	0.038	± 0.0084	0.042	± 0.0068
March, 1996									
Takane-machi, YAMANASHI	6.70	0.998	1.40	0.0069	± 0.0062	0.0069	± 0.0062	0.0031	± 0.0041
Kamiita-machi, TOKUSHIMA	7.80	1.19	1.62	0.018	± 0.0047	0.015	± 0.0040	0.0052	± 0.0045
Koushi-machi, KUMAMOTO	7.31	1.11	1.57	0.020	± 0.0075	0.018	± 0.0067	0.011	± 0.0047

(3)-2

Strontium-90 and Cesium-137 in Milk (producing districts for WHO program)
 (from Oct. 1995 to Mar. 1996)
 -continued from No. 115 of this publication-
 Table (3)-2 Strontium-90 and Cesium-137 in Milk

Location	Component			⁹⁰ Sr				¹³⁷ Cs			
	Ash(g/ℓ)	Ca(g/ℓ)	K(g/ℓ)	(Bq/ℓ)		(Bq/gCa)		(Bq/ℓ)		(Bq/gK)	
November, 1995											
Hokudainoujou, HOKKAIDOU	7.29	1.17	1.62	0.037	± 0.0063	0.031	± 0.0054	0.048	± 0.0071	0.030	± 0.0044
Hachijo-Island, TOKYO	6.94	0.928	1.47	0.061	± 0.010	0.066	± 0.011	0.075	± 0.0088	0.051	± 0.0060
Nishikawa-machi, NIIGATA	7.81	1.10	1.60	0.019	± 0.0042	0.018	± 0.0038	0.014	± 0.0063	0.0086	± 0.0039
Katsuyama, FUKUI	7.34	1.14	1.65	0.024	± 0.0045	0.021	± 0.0039	0.024	± 0.0054	0.015	± 0.0033
Shijounawate, OSAKA	7.28	1.12	1.45	0.043	± 0.0054	0.038	± 0.0048	0.017	± 0.0050	0.011	± 0.0034
Hikawa-machi, SHIMANE	9.96	1.14	1.53	0.055	± 0.0061	0.048	± 0.0053	0.098	± 0.010	0.064	± 0.0067
Takamiya-machi, HIROSHIMA	7.08	1.10	1.54	0.036	± 0.0051	0.032	± 0.0046	0.019	± 0.0052	0.013	± 0.0034
Kochi, KOCHI	7.38	1.16	1.64	0.027	± 0.0045	0.024	± 0.0039	0.013	± 0.0047	0.0080	± 0.0029
Yasu-machi, FUKUOKA	7.22	1.13	1.54	0.023	± 0.0045	0.021	± 0.0040	0.0085	± 0.0043	0.0055	± 0.0028
Kajiki-machi, KAGOSHIMA	7.44	1.17	1.62	0.024	± 0.0047	0.021	± 0.0041	0.023	± 0.0057	0.014	± 0.0035
January, 1996											
Shijounawate, OSAKA	7.14	1.08	1.41	0.039	± 0.0054	0.036	± 0.0050	0.033	± 0.0058	0.024	± 0.0041
Takamiya-machi, HIROSHIMA	7.15	1.07	1.51	0.019	± 0.0066	0.017	± 0.0061	0.015	± 0.0048	0.010	± 0.0032
February, 1996											
Hokudainoujou, HOKKAIDOU	7.40	1.17	1.63	0.032	± 0.0076	0.028	± 0.0065	0.021	± 0.0073	0.013	± 0.0045
Hachijo-Island, TOKYO	6.98	0.990	1.42	0.042	± 0.0058	0.042	± 0.0058	0.022	± 0.0055	0.016	± 0.0039
Nishikawa-machi, NIIGATA	7.71	1.09	1.68	0.021	± 0.0079	0.020	± 0.0072	0.012	± 0.0050	0.0072	± 0.0030
Katsuyama, FUKUI	7.36	1.10	1.75	0.016	± 0.0065	0.014	± 0.0059	0.027	± 0.0069	0.016	± 0.0040
Hikawa-machi, SHIMANE	7.88	1.24	1.65	0.030	± 0.0052	0.024	± 0.0042	0.018	± 0.0055	0.011	± 0.0033
Kochi, KOCHI	7.61	1.21	1.55	0.047	± 0.0090	0.039	± 0.0075	0.0046	± 0.0046	0.0029	± 0.0029
Yasu-machi, FUKUOKA	7.18	1.06	1.52	0.026	± 0.0069	0.024	± 0.0065	0.013	± 0.0055	0.0083	± 0.0036
Kajiki-machi, KAGOSHIMA	7.42	1.14	1.56	0.037	± 0.0086	0.032	± 0.0076	0.021	± 0.0059	0.014	± 0.0038

(3)-3 Strontium-90 and Cesium-137 in Milk (consuming districts)

(from Oct. 1995 to Mar. 1996)

-continued from No. 115 of this publication-

Table (3)-3 Strontium-90 and Cesium-137 in Milk

Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(g/ℓ)	Ca(g/ℓ)	K(g/ℓ)	(Bq/ℓ)	(Bq/gCa)		(Bq/ℓ)	(Bq/gK)	
October, 1995									
Kyoto, KYOTO	7.24	1.09	1.56	0.019 ± 0.0073	0.017 ± 0.0067		0.0093 ± 0.0047	0.0060 ± 0.0030	
December, 1995									
Akita, AKITA	6.03	0.906	1.32	0.026 ± 0.0041	0.028 ± 0.0045		0.013 ± 0.0039	0.0098 ± 0.0029	
January, 1996									
Osaka, OSAKA	7.28	1.10	1.59	0.026 ± 0.0047	0.024 ± 0.0043		0.028 ± 0.0057	0.017 ± 0.0036	
Hiroshima, HIROSHIMA	6.95	1.07	1.49	0.022 ± 0.0069	0.020 ± 0.0064		0.014 ± 0.0046	0.0091 ± 0.0031	
Febraly, 1996									
Sapporo, HOKKAIDOU	7.33	1.11	1.58	0.035 ± 0.0078	0.032 ± 0.0070		0.038 ± 0.0076	0.024 ± 0.0048	
Yamagata, YAMAGATA	6.95	1.06	1.48	0.018 ± 0.0042	0.017 ± 0.0040		0.0088 ± 0.0069	0.0059 ± 0.0046	
Fukushima, FUKUSHIMA	11.2	1.69	2.39	0.031 ± 0.0046	0.018 ± 0.0027		0.023 ± 0.0070	0.0095 ± 0.0029	
Urawa, SAITAMA	7.18	1.09	1.55	0.053 ± 0.0062	0.049 ± 0.0057		0.0057 ± 0.0045	0.0037 ± 0.0029	
Shinjuku, TOKYO	6.98	1.01	1.50	0.016 ± 0.0042	0.016 ± 0.0041		0.0068 ± 0.0046	0.0045 ± 0.0031	
Yokohama, KANAGAWA	7.38	1.11	1.57	0.030 ± 0.0080	0.027 ± 0.0072		0.085 ± 0.0089	0.054 ± 0.0056	
Niigata, NIIGATA	7.44	1.09	1.54	0.030 ± 0.0084	0.027 ± 0.0078		0.017 ± 0.0056	0.011 ± 0.0036	
Fukui, FUKUI	7.42	1.13	1.62	0.025 ± 0.0073	0.022 ± 0.0065		0.023 ± 0.0064	0.014 ± 0.0039	
Nagano, NAGANO	6.50	0.985	1.39	0.026 ± 0.0043	0.026 ± 0.0044		0.0000 ± 0.0037	0.0000 ± 0.0026	
Shizuoka, SHIZUOKA	7.26	1.10	1.57	0.036 ± 0.0054	0.033 ± 0.0049		0.019 ± 0.0056	0.012 ± 0.0036	
Nagoya, AICHI	7.39	1.08	1.52	0.033 ± 0.0062	0.031 ± 0.0058		0.0026 ± 0.0042	0.0017 ± 0.0028	
Yonago, TOTTORI	7.27	1.10	1.56	0.019 ± 0.0045	0.017 ± 0.0041		0.011 ± 0.0049	0.0072 ± 0.0032	
Matsue, SHIMANE	7.67	1.10	1.50	0.026 ± 0.0051	0.024 ± 0.0046		0.014 ± 0.0053	0.0093 ± 0.0035	
Okayama, OKAYAMA	7.20	1.06	1.52	0.021 ± 0.0044	0.020 ± 0.0042		0.012 ± 0.0056	0.0079 ± 0.0037	
Yamaguchi, YAMAGUCHI	7.26	1.07	1.52	0.017 ± 0.0079	0.015 ± 0.0074		0.033 ± 0.0062	0.022 ± 0.0041	
Matsuyama, EHIME	7.17	1.08	1.53	0.029 ± 0.0052	0.027 ± 0.0048		0.0096 ± 0.0047	0.0063 ± 0.0031	
Kochi, KOCHI	7.18	1.11	1.57	0.034 ± 0.0048	0.030 ± 0.0043		0.0090 ± 0.0055	0.0058 ± 0.0035	
Chikushino, FUKUOKA	7.32	1.07	1.57	0.020 ± 0.0067	0.019 ± 0.0063		0.011 ± 0.0060	0.0072 ± 0.0038	
Nagasaki, NAGASAKI	6.89	1.05	1.50	0.024 ± 0.0074	0.022 ± 0.0070		0.018 ± 0.0049	0.012 ± 0.0033	

Location	Component			⁸⁹ Sr			¹³⁷ Cs		
	Ash(g/l)	Ca(g/l)	K(g/l)	(Bq/l)	(Bq/gCa)		(Bq/l)	(Bq/gK)	
Kagoshima, KAGOSHIMA	7.34	1.15	1.55	0.029 ± 0.0081	0.025 ± 0.0070		0.018 ± 0.0056	0.012 ± 0.0036	
Yonagusuku-mura, Okinawa	7.34	1.07	1.54	0.024 ± 0.0048	0.023 ± 0.0045		0.0000 ± 0.0045	0.0000 ± 0.0029	

Table (3)-4 Strontium-90 and Cesium-137 in Milk

Location	Component			⁹⁰ Sr				¹³⁷ Cs			
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kg)		(Bq/gCa)		(Bq/kg)		(Bq/gK)	
January, 1996											
Sample C,	8.00	12.3	17.0	0.36	± 0.021	0.029	± 0.0017	2.0	± 0.05	0.12	± 0.03
February, 1996											
Sample A,	7.92	11.9	17.0	0.44	± 0.024	0.037	± 0.0020	0.34	± 0.022	0.020	± 0.015
Sample B,	2.47	3.11	5.85	0.039	± 0.0086	0.013	± 0.0028	0.14	± 0.011	0.023	± 0.008
Sample D,	2.58	3.92	5.81	0.038	± 0.0085	0.0096	± 0.0022	0.032	± 0.0065	0.0055	± 0.002
Sample E,	2.39	3.68	5.26	0.069	± 0.0069	0.019	± 0.0019	0.10	± 0.010	0.019	± 0.006
Sample F,	2.53	3.16	4.98	0.062	± 0.010	0.020	± 0.0033	0.14	± 0.011	0.028	± 0.009

(4)-1 Strontium-90 and cesium-137 in Vegetables (producing districts)
(from Oct. 1995 to Mar. 1996)

-continued from No. 115 of this publication-

Table (4)-1 :Strontium-90 and cesium-137 in Vegetables

Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)		(Bq/gCa)	(Bq/kgwet)		(Bq/gK)
<u>(Cabbage)</u>									
October, 1995									
Mutsu, AOMORI	0.586	0.399	2.15	0.15	± 0.010	0.39	± 0.024	0.045	± 0.0087
November, 1995	0.527	0.437	1.95	0.21	± 0.017	0.48	± 0.038	0.028	± 0.0070
Sannohe-machi, AOMORI									
January, 1996									
Kumatori-machi, OSAKA	0.625	0.293	2.48	0.031	± 0.0050	0.11	± 0.017	0.0008	± 0.0039
<u>(Chinese cabbage)</u>									
October, 1995									
Tamayama-mura, IWATE	0.546	0.431	2.21	0.11	± 0.008	0.26	± 0.019	0.016	± 0.0053
December, 1995									
Utsunomiya, TOCHIGI	0.595	0.415	2.31	0.26	± 0.019	0.63	± 0.045	0.092	± 0.011
January, 1996									
Shinguu, WAKAYAMA	0.547	0.412	2.03	0.31	± 0.014	0.75	± 0.033	0.0055	± 0.0057
<u>(Japanese radish)</u>									
October, 1995									
Tamayama-mura, IWATE	0.617	0.296	2.71	0.042	± 0.0054	0.14	± 0.018	0.011	± 0.0046
Uno-ke-machi, ISHIKAWA	0.534	0.103	2.25	0.16	± 0.010	1.6	± 0.10	0.0097	± 0.0053
Takamatsu, KAGAWA	0.362	0.155	1.32	0.14	± 0.009	0.90	± 0.055	0.037	± 0.0055
November, 1995									
Sannohe-machi, AOMORI	0.475	0.247	1.91	0.26	± 0.018	1.0	± 0.07	0.027	± 0.0066
Fukushima, FUKUSHIMA	0.600	0.264	2.27	0.035	± 0.0052	0.13	± 0.020	0.0007	± 0.0032
Mito, IBARAKI	0.512	0.305	1.94	0.082	± 0.0069	0.27	± 0.022	0.0072	± 0.0055
Maebashi, GUNMA	0.731	0.237	3.10	0.018	± 0.0043	0.076	± 0.018	0.0008	± 0.0036
Kosugi-machi, TOYAMA	0.364	0.153	1.50	0.024	± 0.0054	0.16	± 0.035	0.0008	± 0.0057
Fukui, FUKUI	0.527	0.163	1.86	0.0077	± 0.0033	0.048	± 0.020	0.0011	± 0.0049

Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)	(Bq/gCa)		(Bq/kgwet)	(Bq/gK)	
Toyama, TOYAMA	1.63	0.927	6.76	0.050 ± 0.0067	0.054 ± 0.0072		0.012 ± 0.0068	0.0018 ± 0.0010	
Kusu-machi, MIE	1.46	0.701	5.98	0.23 ± 0.012	0.33 ± 0.017		0.013 ± 0.0078	0.0022 ± 0.0013	
Takamatsu, KAGAWA	1.54	0.612	6.30	0.028 ± 0.0049	0.046 ± 0.0081		0.0078 ± 0.0044	0.0012 ± 0.00069	
Matsumoto-machi, KAGOSHIMA	1.43	0.603	4.28	0.040 ± 0.0055	0.067 ± 0.0091		0.11 ± 0.010	0.025 ± 0.0023	
November, 1995									
Fukushima, FUKUSHIMA	1.86	1.15	6.32	0.11 ± 0.008	0.094 ± 0.0070		0.039 ± 0.0069	0.0061 ± 0.0011	
Mito, IBARAKI	1.65	0.949	6.81	0.14 ± 0.009	0.15 ± 0.009		0.058 ± 0.0086	0.0085 ± 0.0013	
Maebashi, GUNMA	2.13	0.511	9.02	0.031 ± 0.0051	0.061 ± 0.0099		0.0022 ± 0.0039	0.00024 ± 0.00043	
Fukui, FUKUI	1.56	0.350	6.76	0.064 ± 0.0079	0.18 ± 0.023		0.0022 ± 0.0057	0.00032 ± 0.00084	
Saku, NAGANO	1.84	0.684	7.75	0.035 ± 0.0058	0.051 ± 0.0085		0.0050 ± 0.0042	0.00064 ± 0.00054	
Gifu, GIFU	1.65	0.769	6.38	0.047 ± 0.0061	0.061 ± 0.0079		0.012 ± 0.0054	0.0018 ± 0.00084	
Gotenba, SHIZUOKA	1.65	1.17	5.48	0.26 ± 0.013	0.22 ± 0.011		0.49 ± 0.021	0.089 ± 0.0038	
Rittou-machi, SHIGA	1.94	0.954	8.35	0.11 ± 0.008	0.12 ± 0.009		0.0000 ± 0.0040	0.00000 ± 0.00048	
Kasai, HYOUGO	1.64	0.597	6.47	0.093 ± 0.0085	0.16 ± 0.014		0.011 ± 0.0047	0.0017 ± 0.00073	
Kurayoshi, TOTTORI	1.70	0.762	6.96	0.098 ± 0.0084	0.13 ± 0.011		0.034 ± 0.0071	0.0049 ± 0.0010	
Matsuyama, EHIME	1.95	0.936	7.70	0.053 ± 0.0062	0.057 ± 0.0066		0.016 ± 0.0066	0.0020 ± 0.00085	
Shime-machi, FUKUOKA	1.77	0.831	7.56	0.028 ± 0.0050	0.034 ± 0.0060		0.0077 ± 0.0064	0.0010 ± 0.00085	
Saga, SAGA	1.78	0.673	7.82	0.038 ± 0.0053	0.056 ± 0.0079		0.0017 ± 0.0038	0.00022 ± 0.00049	
Takanabe-machi, MIYAZAKI	1.69	0.525	7.07	0.28 ± 0.013	0.53 ± 0.024		0.0004 ± 0.0041	0.00006 ± 0.00058	
December, 1995									
Chiba, CHIBA	1.93	0.301	7.01	0.032 ± 0.0083	0.11 ± 0.028		0.0000 ± 0.0055	0.00000 ± 0.00079	
Takane-machi, YAMANASHI	3.08	1.48	11.7	0.13 ± 0.009	0.089 ± 0.0061		0.0081 ± 0.0050	0.00070 ± 0.00042	
Kashihara, NARA	1.98	0.469	8.83	0.0000 ± 0.0033	0.0000 ± 0.0070		0.0000 ± 0.0044	0.00000 ± 0.00050	
Kubokawa-machi, KOCHI	2.10	0.817	8.36	0.13 ± 0.009	0.16 ± 0.011		0.029 ± 0.0072	0.0035 ± 0.00086	
Usa, OITA	1.74	0.304	7.85	0.062 ± 0.0069	0.20 ± 0.023		0.0000 ± 0.0038	0.00000 ± 0.00048	
January, 1996									

Location	Component			⁹⁰ Sr				¹³⁷ Cs			
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kg wet)		(Bq/g Ca)		(Bq/kg wet)		(Bq/g K)	
Hirosima, HIROSHIMA	1.45	0.321	5.70	0.021	± 0.0049	0.067	± 0.015	0.0087	± 0.0041	0.0015	± 0.00072
Yuya-machi, YAMAGUCHI	1.82	0.568	7.37	0.23	± 0.012	0.41	± 0.021	0.018	± 0.0072	0.0024	± 0.00098
Ishii-machi, TOKUSHIMA	1.95	0.809	6.86	0.12	± 0.009	0.14	± 0.011	0.036	± 0.0080	0.0052	± 0.0012

(4)-2 Strontium-90 and cesium-137 in Vegetables (consuming districts)
(from Oct. 1995 to Mar. 1996)

-continued from No. 115 of this publication-

Table (4)-2 :Strontium-90 and cesium-137 in Vegetables

Location	Component			⁹⁰ Sr				¹³⁷ Cs			
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)		(Bq/gCa)		(Bq/kgwet)		(Bq/gK)	
<u>(Cabbage)</u>											
November, 1995											
Akita, AKITA	0.604	0.407	2.35	0.16	± 0.015	0.40	± 0.037	0.014	± 0.0067	0.0062	± 0.0028
<u>(Japanese radish)</u>											
October, 1995											
Yamagata, YAMAGATA	0.462	0.234	1.78	0.0045	± 0.0036	0.019	± 0.015	0.0036	± 0.0039	0.0020	± 0.0022
Kyoto, KYOTO	0.415	0.230	1.48	0.16	± 0.010	0.71	± 0.042	0.016	± 0.0054	0.011	± 0.0037
November, 1995											
Akita, AKITA	0.475	0.239	2.05	0.040	± 0.0083	0.17	± 0.035	0.0009	± 0.0062	0.0004	± 0.0030
Shinjuku, TOKYO	0.537	0.251	2.18	0.061	± 0.0065	0.24	± 0.026	0.023	± 0.0062	0.011	± 0.0028
Niigata, NIIGATA	0.416	0.189	1.56	0.028	± 0.0051	0.15	± 0.027	0.017	± 0.0064	0.011	± 0.0041
Osaka, OSAKA	0.380	0.188	1.30	0.0095	± 0.0039	0.051	± 0.021	0.057	± 0.0086	0.044	± 0.0066
Okayama, OKAYAMA	0.452	0.296	1.74	0.22	± 0.012	0.76	± 0.039	0.014	± 0.0050	0.0080	± 0.0029
December, 1995											
Yonagusuku-mura, Okinawa	0.587	0.243	2.28	0.022	± 0.0048	0.091	± 0.020	0.0004	± 0.0041	0.0002	± 0.0018
January, 1996											
Nagasaki, NAGASAKI	0.512	0.185	1.52	0.061	± 0.0062	0.33	± 0.033	0.016	± 0.0067	0.010	± 0.0044
February, 1996											
Yokohama, KANAGAWA	0.485	0.242	2.00	0.018	± 0.0047	0.073	± 0.020	0.0027	± 0.0052	0.0013	± 0.0026
<u>(Spinach)</u>											
October, 1995											
Yamagata, YAMAGATA	1.55	0.552	6.46	0.12	± 0.008	0.23	± 0.015	0.010	± 0.0045	0.0016	± 0.00069
November, 1995											
Shinjuku, TOKYO	1.96	0.368	9.09	0.029	± 0.0051	0.078	± 0.014	0.015	± 0.0064	0.0017	± 0.00070
Kyoto, KYOTO	1.45	1.11	5.47	0.046	± 0.0093	0.042	± 0.0084	0.028	± 0.0072	0.0051	± 0.0013
Osaka, OSAKA	1.89	1.02	7.39	0.055	± 0.0062	0.054	± 0.0061	0.029	± 0.0065	0.0039	± 0.00087

Location	Component			⁹⁰ Sr				¹³⁷ Cs			
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)		(Bq/gCa)		(Bq/kgwet)		(Bq/gK)	
Okayama, OKAYAMA	1.94	0.443	8.28	0.013	± 0.0043	0.030	± 0.0097	0.013	± 0.0051	0.0016	± 0.00061
Matsuyama, EHIME	1.76	0.319	7.48	0.026	± 0.0049	0.082	± 0.015	0.0077	± 0.0047	0.0010	± 0.00062
December, 1995											
Yonagusuku-mura, Okinawa	1.59	0.536	6.73	0.011	± 0.0042	0.020	± 0.0078	0.0000	± 0.0058	0.00000	± 0.00087
January, 1996											
Nagasaki, NAGASAKI	1.64	0.491	6.78	0.036	± 0.0052	0.074	± 0.011	0.013	± 0.0064	0.0019	± 0.00094
February, 1996											
Yokohama, KANAGAWA	1.96	0.204	9.05	0.035	± 0.0057	0.17	± 0.028	0.017	± 0.0061	0.0019	± 0.00067

(5) Strontium-90 and cesium-137 in Sea Fish
(from Oct. 1995 to Mar. 1996)

-continued from No. 115 of this publication-

Table (5) :Strontium-90 and cesium-137 in Sea Fish

Location	Component			⁹⁰ Sr		¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)	(Bq/gCa)	(Bq/kgwet)	(Bq/gK)	
<u>(Branchiostegus sp)</u>								
November, 1995								
Nagasaki, NAGASAKI	1.10	0.378	3.63	0.0000 ± 0.0027	0.0000 ± 0.0072	0.18 ± 0.012	0.049 ± 0.0033	
<u>(Limanda herzensteini)</u>								
November, 1995								
Mutsu, AOMORI	1.35	0.845	3.70	0.0049 ± 0.0059	0.0058 ± 0.0070	0.095 ± 0.010	0.026 ± 0.0027	
Niigata, NIIGATA	1.40	0.764	3.95	0.0006 ± 0.0032	0.0008 ± 0.0041	0.094 ± 0.011	0.024 ± 0.0027	
Mikuni-machi, FUKUI	2.44	5.23	3.30	0.0037 ± 0.0057	0.0007 ± 0.0011	0.13 ± 0.011	0.040 ± 0.0034	
Aji-machi, KAGAWA	3.44	7.40	3.44	0.013 ± 0.0040	0.0017 ± 0.00055	0.073 ± 0.010	0.021 ± 0.0029	
February, 1996								
Ootake, HIROSHIMA	1.72	2.37	3.46	0.0031 ± 0.0048	0.0013 ± 0.0020	0.056 ± 0.0082	0.016 ± 0.0024	
<u>(Mugil cephalus)</u>								
November, 1995								
Ushimado-machi, OKAYAMA	1.39	0.305	4.40	0.0022 ± 0.0030	0.007 ± 0.010	0.087 ± 0.0097	0.020 ± 0.0022	
<u>(Pterocaesio diagramma)</u>								
December, 1995								
Yonagusuku-mura, Okinawa	3.49	8.21	4.43	0.015 ± 0.0040	0.0018 ± 0.00049	0.15 ± 0.013	0.034 ± 0.0029	
<u>(Sardinops melanostictus)</u>								
February, 1996								
Nagano, NAGANO	3.04	6.59	3.08	0.0049 ± 0.0030	0.00075 ± 0.00046	0.057 ± 0.0091	0.018 ± 0.0030	
<u>(Scomber australasicus)</u>								
February, 1996								
Chikura-machi, CHIBA	0.998	0.177	3.01	0.0006 ± 0.0030	0.003 ± 0.017	0.13 ± 0.012	0.043 ± 0.0039	
<u>(Scomber sp)</u>								
November, 1995								
Kyoto, KYOTO	1.17	0.228	3.55	0.0028 ± 0.0039	0.012 ± 0.017	0.11 ± 0.011	0.031 ± 0.0032	

Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)	(Bq/gCa)		(Bq/kgwet)	(Bq/gK)	
Osaka, OSAKA January, 1996	1.10	0.102	3.20	0.0000 ± 0.0032	0.000 ± 0.031		0.12 ± 0.010	0.038 ± 0.0032	
Oki-adjacent seas, TOTTRI (<i>Sebastes inermis</i>) January, 1996	1.39	0.264	3.29	0.0000 ± 0.0047	0.000 ± 0.018		0.16 ± 0.013	0.048 ± 0.0040	
Yamaguchi, YAMAGUCHI (<i>Seriola quinqueradiata</i>) October, 1995	5.18	14.5	3.30	0.024 ± 0.0047	0.0016 ± 0.00032		0.15 ± 0.013	0.046 ± 0.0039	
Togi-machi, ISHIKAWA (<i>Spratelloides gracilis</i>) November, 1995	0.694	0.409	2.14	0.0031 ± 0.0054	0.008 ± 0.013		0.12 ± 0.011	0.056 ± 0.0050	
Akune, KAGOSHIMA (<i>Trachurus sp.</i>) November, 1995	2.65	5.50	3.28	0.0058 ± 0.0038	0.0011 ± 0.00068		0.15 ± 0.012	0.046 ± 0.0037	
Miyake-island, TOKYO	1.26	1.54	2.69	0.0052 ± 0.0035	0.0034 ± 0.0023		0.12 ± 0.011	0.044 ± 0.0042	
Odawara, KANAGAWA	1.62	0.921	4.47	0.0000 ± 0.0032	0.0000 ± 0.0034		0.17 ± 0.013	0.039 ± 0.0028	
Shizuoka, SHIZUOKA	3.25	7.82	3.18	0.022 ± 0.0044	0.0028 ± 0.00056		0.19 ± 0.013	0.059 ± 0.0042	

Sea Fish

Japanese name	English name	Scientific name
Amadai	Tilefish	<u>Branchiostegus</u> sp
Magarei	Brown sole	<u>Limanda herzensteini</u>
Bora	Gray mullet	<u>Mugil cephalus</u>
Takasago	Golden banded fusilier	<u>Pterocaesio diagramma</u>
Maiwashi	Japanese pilchard	<u>Sardinops melanostictus</u>
Gomasaba	Spotted chub mackerel	<u>Scomber australasicus</u>
Saba	Mackerel	<u>Scomber</u> sp
Mebaru	Black rockfish	<u>Sebastes inermis</u>
Buri	Yellow-tail	<u>Seriola quinqueradiata</u>
Kibinago	Blue sprat	<u>Spratelloides gracilis</u>
Aji	Horse mackerel	<u>Trachurus</u> sp

(6) Strontium-90 and cesium-137 in Freshwater Fish
 (from Oct. 1995 to Mar. 1996)

-continued from No. 115 of this publication-

Table (6) :Strontium-90 and cesium-137 in Freshwater Fish

Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)	(Bq/gCa)		(Bq/kgwet)	(Bq/gK)	
<u>(Carassius auratus)</u>									
November, 1995									
Niigata, NIIGATA	1.10	0.720	3.12	0.077	± 0.011	0.11	± 0.015	0.18	± 0.018
December, 1995									
Mikata-machi, FUKUI	1.49	2.13	3.28	0.18	± 0.010	0.083	± 0.0048	0.20	± 0.014
Uji, KYOTO	4.47	13.2	2.79	0.67	± 0.021	0.051	± 0.0016	0.018	± 0.0072
<u>(Cyprinus carpio)</u>									
October, 1995									
Shobara, HIROSHIMA	1.00	0.411	3.23	0.043	± 0.0056	0.11	± 0.014	0.075	± 0.0084
<u>(Hypomesus nipponensis)</u>									
December, 1995									
Suwa, NAGANO	2.21	5.13	2.58	0.082	± 0.0071	0.016	± 0.0014	0.11	± 0.010
<u>(Salmo gairdneri)</u>									
November, 1995									
Kumagaya, SAITAMA	1.15	0.139	4.13	0.0003	± 0.0032	0.002	± 0.023	0.12	± 0.010
								0.028	± 0.0025

Freshwater Fish

Japanese name	English name	Scientific name
Funa	Crucian carp	<u>Carassius auratus</u>
Koi	Carp	<u>Cyprinus carpio</u>
Wakasagi	Japanese smelt	<u>Hypomesus japonensis</u>
Nijimasu	Rainbow trout	<u>Salmo gairdneri</u>

(7) Strontium-90 and cesium-137 in Shellfish
 (from Oct. 1995 to Mar. 1996)

-continued from No. 115 of this publication-

Table (7) :Strontium-90 and cesium-137 in Shellfish

Location	Component			⁹⁰ Sr		¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)	(Bq/gCa)	(Bq/kgwet)	(Bq/gK)	
<u>(<i>Patinopecten yessoensis</i>)</u>								
November, 1995								
Mutsu, AOMORI	1.46	0.217	2.37	0.0000 ± 0.0047	0.000 ± 0.022	0.022 ± 0.0072	0.0094 ± 0.0030	
Febraly, 1996								
Yamada-machi, IWATE	2.47	0.306	3.83	0.0033 ± 0.0036	0.011 ± 0.012	0.039 ± 0.0082	0.010 ± 0.0021	

Shellfish

Japanese name	English name	Scientific name
Hotategai	Yesso scallop	<u><i>Patinopecten yessoensis</i></u>

(30)

(8) Strontium-90 and cesium-137 in Seaweeds
 (from Oct. 1995 to Mar. 1996)

-continued from No. 115 of this publication-

Table (8) :Strontium-90 and cesium-137 in Seaweeds


Location	Component			⁹⁰ Sr			¹³⁷ Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	(Bq/kgwet)	(Bq/gCa)		(Bq/kgwet)	(Bq/gK)	
<u>(<i>Undaria pinnatifida</i>)</u>									
February, 1996									
Minamichita-machi, AICHI	1.97	0.727	5.64	0.046 ± 0.0091	0.063 ± 0.013		0.036 ± 0.0076	0.0064 ± 0.0014	
Hiroshima, HIROSHIMA	1.73	0.502	5.54	0.024 ± 0.0044	0.048 ± 0.0088		0.0088 ± 0.0063	0.0016 ± 0.0011	
Shimabara, NAGASAKI	2.08	0.525	6.87	0.0095 ± 0.0037	0.018 ± 0.0071		0.014 ± 0.0072	0.0020 ± 0.0010	

Seaweeds

Japanese name	English name	Scientific name
Wakame	Wakame seaweed	<u>Undaria pinnatifida</u>

* * Total Diet * *

<Strontium-90>



<Cesium-137>





Fig. 1

* * Rice (producing districts) * *

<Strontium-90>



<Cesium-137>





Fig.2-1

* * Rice (consuming districts) * *

<Strontium-90>



<Cesium-137>





Fig.2-2

* * Milk (producing districts for domestic program)

<Strontium-90>



<Cesium-137>





Fig.3-1

* * Milk (producing districts for WHO program) * *

<Strontium-90>



<Cesium-137>





Fig.3-2

* * Milk (consuming districts) * *

<Strontium-90>



<Cesium-137>





Fig.3-3

* * Powdered Milk * *

<Strontium-90>



<Cesium-137>





Fig. 3-4

* * Vegetables (producing districts) * *

<Strontium-90>



<Cesium-137>





Fig.4-1

* * Vegetables (consuming districts) * *

<Strontium-90>



<Cesium-137>





Fig. 4-2

* * Sea Fish * *

<Strontium-90>



<Cesium-137>





Fig. 5

* * Freshwater Fish * *

<Strontium-90>



<Cesium-137>





Fig. 6

* * Shellfish * *

<Strontium-90>



<Cesium-137>





Fig. 7

* * Seaweeds * *

<Strontium-90>



<Cesium-137>




Fig. 8

* * Sampling Locations in Japan * *

- | | |
|----------------|---------------|
| 1: Sapporo | 36: Hiroshima |
| 2: Aomori | 37: Kochi |
| 3: Morioka | 38: Matsuyama |
| 4: Akita | 39: Yamaguchi |
| 5: Sendai | 40: Ooita |
| 6: Yamagata | 41: Fukuoka |
| 7: Fukushima | 42: Saga |
| 8: Niigata | 43: Kumamoto |
| 9: Mito | 44: Miyazaki |
| 10: Utsunomiya | 45: Nagasaki |
| 11: Chiba | 46: Kagoshima |
| 12: Urawa | 47: Naha |
| 13: Shinjuku | |
| 14: Maebashi | |
| 15: Nagano | |
| 16: Yokohama | |
| 17: Toyama | |
| 18: Kouhu | |
| 19: Kanazawa | |
| 20: Shizuoka | |
| 21: Gifu | |
| 22: Fukui | |
| 23: Nagoya | |
| 24: Tsu | |
| 25: Ootsu | |
| 26: Kyoto | |
| 27: Nara | |
| 28: Osaka | |
| 29: Tottori | |
| 30: Kobe | |
| 31: Wakayama | |
| 32: Okayama | |
| 33: Matsue | |
| 34: Tokushima | |
| 35: Takamatsu | |

