



ISSN 0441-2516

NIRS-RSD-122

RADIOACTIVITY SURVEY DATA in Japan

Part 1
= Environmental Materials =

NUMBER 122
August 1999

National Institute of Radiological Sciences
Chiba, Japan

Radioactivity Survey Data
in Japan
Number 122

August 1999 part 1 = Environmental Materials =

Contents

	Page
Environmental and Dietary Materials	
(Japan Chemical Analysis Center)	
1. Collection and pretreatment of samples	1
2. Preparation of samples for analysis	3
3. Separation of Strontium-90 and Cesium-137	3
4. Determination of Stable Strontium, Calcium and Potassium	4
5. Counting	4
6. Results	5
(1) -1 Strontium-90 and Cesium-137 in Rain and Dry Fallout	5
(for domestic program)	
-2 Strontium-90 and Cesium-137 in Rain and Dry Fallout	14
(for WHO program)	
(2) Strontium-90 and Cesium-137 in Airborne Dust	17
(3) Strontium-90 and Cesium-137 in Service Water	20
(4) Strontium-90 and Cesium-137 in Freshwater	23
(5) Strontium-90 and Cesium-137 in Soil	24
(6) Strontium-90 and Cesium-137 in Sea Water	28
(7) Strontium-90 and Cesium-137 in Sea Sediments	29
7. Contents of Figure (Selected Locations)	30

Edited by National Institute of Radiological Sciences, under the supervision of Science and Technology Agency of Japanese Government.

Environmental and Dietary Materials*

(Japan Chemical Analysis Center)

1. Collection and pretreatment of samples

(1) Rain and dry fallout

Rain and dry fallout was collected monthly on a sampling tray, approximately 5000cm² in area, which was filled with water to a depth of 1 cm at the beginning of every month.

Strontium and cesium carrier solutions were added after the sample was filtered. The tray was washed with 5 ℓ of distilled water and the washing was combined to the filtrate.

The sample was passed through a cation exchange column (500m ℓ of Dowex 50W X8, 50~100 mesh, Na form) at a rate flow of 80m ℓ /min.

(2) Airborne dust

Airborne dust was collected by an electrostatic precipitator or a filter air sampler for every three-months at a rate of more than 3000m³ per month.

The sampling was done 1 to 1.5 meters above the ground.

(3) Service water and freshwater

Service water, 100 ℓ each, was collected at the intake of the water-treatment plant and at the tap after water was left running for five minutes. Strontium and cesium carriers were added to the filtered water sample. The subsequent process was the same as that described in the section (1). Freshwater was treated in the same way as the service water.

(4) Soil

Soil was collected from the location in the spacious and flat area without past surface disturbance caused by dust storms, inflow and out flow due to precipitation, etc.. Any places located under trees in a forest, in a stony area or inside of river banks were avoided. Soil was taken from two layers of different depths, 0-5cm and 5-20cm. The soil lumps were crushed by hands and dried in a drying oven regulated 105°C . The soil was then passed through a 2mm sieve to remove plant roots and pebbles.

(5) Sea water

Sea water was collected at the fixed stations

where the effect of terrestrial fresh water from rivers was expected to be negligibly small. A special consideration was also given to weather conditions.

The sampling was carried out when there was no rainfall for the last few days. To prevent contamination, water samples were collected at the bow of a sampling boat just before she stood still by scooping surface water using a polyethylene bucket.

Immediately after the collection, the samples were acidified to a pH lower than 3 by adding concentrated hydrochloric acid in a ratio of 1m ℓ to 1 ℓ of sea water, and then stored in 20 ℓ polyethylene containers. The sampling equipments as well as containers were thoroughly rinsed with dilute hydrochloric acid and then with distilled water before use. Two hundred milliliters of sea water was also collected at the same stations for the determination of chlorinity.

(6) Sea sediments

Sediment was collected in the same area as that for the sea water sample, taking the following criteria into account:

- a. The depth of water exceeds 1m at low tide.
- b. No significant sedimental movement is observed in the vicinity of concern.
- c. Mud, silt and fine sand are preferable.

A conventional sediment sampling device was used for collecting the top few centimeters of surface sediment. Approximately 4 kg of the sample in wet weight was spread on a stainless steel dish after removed of the pebbles, shells and other foreign materials, and dried in a drying oven regulated at 105°C.

(7) Total diet

A full one day ordinary diet including three meals, water, tea and other in-between snacks for five persons was collected as a sample of "total diet".

The sample in a large stainless steel pan was carbonized carefully by direct application of gas flame, and was transferred to a porcelain dish and then ashed at 450°C in an electric muffle furnace.

(8) Rice

Polished rice was collected in producing districts at the harvest and in consuming areas when new crops were first put on sale. The sample was carbonized and ashed in a porcelain dish.

* Samples were sent to the Center from 46 contracted prefectures.

(9) Milk

Raw milk was collected in producing districts and commercial milk was purchased in consuming districts. Milk in a stainless steel pan or a porcelain dish was evaporated to dryness followed by carbonization and ashing.

(10) Vegetables

Spinach and Japanese radish were selected as the representatives for leaf vegetables and for nonstarch roots, respectively. After removing soil, the edible part of vegetable sample was dried and carbonized in a stainless steel pan or a porcelain dish.

(11) Tea

Five hundred grams of manufactured green tea was collected, carbonized and ashed in a stainless steel pan or a porcelain dish.

(12) Fish, shellfish and seaweeds

a. Sea fish and freshwater fish

Fish was rinsed with water and blotted with a filter paper. Only the edible part was used in case of larger sized fish, and the whole part was used in case of smaller ones. Each sample was weighed and placed in a stainless steel pan or a porcelain dish. After carbonized, the sample was ashed in an electric muffle furnace.

b. Shellfish

Approximately 4kg of shellfish including the shells was collected or purchased. After removing the shells, it was treated in the same way as that for the sea fish.

c. Seaweeds

Edible seaweeds were collected and rinsed with water to remove sand and other adhering matters on the surface. These were removed of excess water, weighed dried and ashed.

Table 1 shows details of sample collection.

Table 1 Details of sample collection

Sample	Frequency of sampling	Quantity of sample
=Environmental materials=		
(1) Rain and dry fallout		
1. For domestic program	monthly	
2. For WHO program	monthly	
(2) Airborne dust	quarterly	>3000 m ³ /month
(3) Service water and freshwater		
1. Service water (source water)	semiyearly	100 ℥
2. Service water (tap water)	semiyearly	100 ℥
3. Freshwater	yearly (fishing season)	100 ℥
(4) Soil		
1. 0~ 5 cm	yearly	4 kg
2. 5~ 20cm	yearly	4 kg
(5) Sea water	yearly	40 ℥
(6) Sea sediments	yearly	4 kg
=Dietary materials=		
(7) Total diet	semiyearly	daily amount for 5 persons
(8) Rice		
1. Producing districts	yearly (harvesting season)	5 kg (polished rice)
2. Consuming districts	yearly (harvesting season)	5 kg (polished rice)
(9) Milk		
1. Producing districts for WHO program	quarterly (February, May, August and November)	3 ℥
2. Producing districts for domestic program	semiyearly (February and August)	3 ℥

Sample	Frequency of sampling	Quantity of sample
3. Consuming districts	semiyearly (February and August)	3 ℥
4. Powdered milk	semiyearly (April and October)	2~ 3 kg
(10) Vegetables		
1. Producing districts	yearly (harvesting season)	4 kg
2. Consuming districts	yearly (harvesting season)	4 kg
(11) Tea	yearly (the first harvesting season)	500g (manufactured tea)
(12) Fish, shellfish and seaweeds		
1. Sea fish	yearly (fishing season)	4 kg
2. Freshwater fish	yearly (fishing season)	4 kg
3. Shellfish	yearly (fishing season)	4 kg
4. Seaweeds	yearly (fishing season)	2~ 3 kg

2. Preparation of samples for analysis

(1) Rain, service water and freshwater

Strontium and cesium were eluted with hydrochloric acid from the cation exchange column. The residue of rain sample on the filter paper was ashed in an electric muffle furnace and the ash was dissolved in hydrochloric acid. The insoluble part was filtered and washed. The filtrate and the washings were combined to the previous eluate and used for radiochemical analysis.

(2) Soil and Sea sediment

Dried soil was crushed to smaller ones than 0.2 5mm in size by a crusher. The sieved sample was ashed in an electric muffle furnace regulated at 450 °C. The sample was then heated with hydrochloric acid, strontium and cesium carrier solutions and the mixture was heated. The insoluble constituent was filtered off and washed with water.

The dried sample was crushed to smaller ones than 0.25mm by a crushing machine. The further preparation of the sample was the same as that described in the section 2-(2).

(3) Rice

The ashed sample was pulverized with a porcelain mortar and passed through a 0.35mm sieve. The sieved sample to which both strontium and cesium carriers were added, was digested with nitric acid by heating. After the sample was heated again with nitric acid to dryness, strontium and cesium were extracted with hydrochloric acid and water. The insoluble constituent was filtered and washed. The filtrate and washings were combined for subsequent radiochemical analysis.

(4) Airborne dust, diet, milk, vegetables, fish and shellfish, seaweeds, tea and others

These ashed samples were treated with the

same procedure as that described in the section 2-(4).

3. Separation of strontium-90 and cesium-137

(1) Strontium-90

Sample solutions, prepared as in the foregoing sections 2-(1) through 2-(4), were neutralized with sodium hydroxide. After sodium carbonate was added, the precipitate of strontium and calcium carbonates was separated. The supernatant solution was retained for cesium-137 determination.

The carbonates were dissolved in hydrochloric acid and strontium and calcium were precipitated as oxalates. The precipitate was dissolved in nitric acid and strontium was separated from calcium by successive fuming nitric acid separation. Iron scavenge was made after addition of ferric iron carrier followed by barium chromate separation after addition of barium carrier to remove radium, its daughters and lead. Strontium was recovered as carbonate, and the precipitate was dried and weighed to determine strontium recovery. The strontium carbonate was dissolved in hydrochloric acid and iron carrier was added. The solution was allowed to stand for two weeks for strontium-90 and yttrium-90 to attain equilibrium. Yttrium-90 was coprecipitated with ferric hydroxide and the precipitate was filtered off, washed and counted.

(2) Cesium-137

The supernatant separated from the strontium fraction was acidified with hydrochloric acid. While stirring, cesium was adsorbed on the ammonium molybdate added.

After filtered off and washed with hydrochloric acid the precipitate was dissolved in 2.5N sodium hydroxide solution. The solution was adjusted to pH 8.2 with hydrochloric acid and allowed to cool.

Resultant molybdenum hydroxide which separated

out in the solution, was filtered off and washed with water. EDTA was added to the filtrate and washings. Cesium and rubidium were adsorbed on a cation exchange column and cesium was separated from rubidium by eluting with hydrochloric acid.

The eluate was evaporated to dryness and was dissolved. The solution was filtered.

Chloroplatinic acid was added to precipitate cesium. The precipitate was filtered onto a tared paper using a demountable filter and washed with water and then ethanol. After drying, the chemical yield of cesium was determined by weighing the precipitate. Cesium-137 radioactivity was measured for this precipitate.

4. Determination of stable strontium, calcium and potassium

A weighed amount of soil or sea sediment was heated in a electric muffle furnace at 450°C and then treated with hydrochloric acid for extraction. A weighed aliquot of ashed samples of total diet, vegetables, milk, fish, shellfish or seaweeds was

digested with hydrofluoric acid and nitric acid. The extract was made up to an appropriate volume with dilute hydrochloric acid. The sample solution was analyzed for calcium by titration with standard potassium permanganate solution after separating calcium as oxalate. Atomic absorption spectroscopy was applied when appropriate. Stable strontium and potassium were determined by atomic absorption and flame emission spectrometry, respectively.

5. Counting

After the radiochemical separation the mounted precipitates were counted for activity using low background beta counters normally for 60 to 90min. Net sample counting rates were corrected for counter efficiency, recovery, self-absorption and decay to obtain the content of strontium-90 and cesium-137 per sample aliquot. From the results, concentrations of these nuclides in the original samples were calculated.

6. Results

(1)-1 Strontium-90 and Cesium-137 in Rain and Dry Fallout(for domestic program)
(from Apr. 1997 to Sep. 1997)

-continued from No. 120 of this publication-

Table (1)-1 : Strontium-90 and Cesium-137 in Rain and Dry Fallout

Location	Duration (days)	Precipitation (mm)	^{90}Sr		^{137}Cs	
			(MBq/km ²)	(MBq/km ²)	(MBq/km ²)	(MBq/km ²)
April, 1997						
Sapporo, HOKKAIDOU	31	10.5	0.019	± 0.0072	0.016	± 0.013
Morioka, IWATE	31	64.3	0.078	± 0.019	0.061	± 0.015
Onagawa-machi, MIYAGI	31	67.5	0.027	± 0.0094	0.002	± 0.010
Yamagata, YAMAGATA	31	74.6	0.055	± 0.015	0.024	± 0.012
Ookuma-machi, FUKUSHIMA	31	58.9	0.044	± 0.014	0.016	± 0.011
Mito, IBARAKI	31	78.0	0.032	± 0.0080	0.036	± 0.014
Kawachi-machi, TOCHIGI	32	132.7	0.040	± 0.0084	0.018	± 0.012
Maebashi, GUNMA	31	52.0	0.014	± 0.0077	0.033	± 0.014
Urawa, SAITAMA	31	91.4	0.016	± 0.0081	0.059	± 0.014
Ichihara, CHIBA	31	124.1	0.029	± 0.0089	0.003	± 0.012
Shinjuku, TOKYO	31	135.1	0.016	± 0.014	0.001	± 0.012
Yokohama, KANAGAWA	32	148.2	0.022	± 0.0073	0.044	± 0.015
Kosugi-machi, TOYAMA	31	247.0	0.037	± 0.0084	0.004	± 0.011
Fukui, FUKUI	31	165.4	0.075	± 0.035	0.000	± 0.054
Koufu, YAMANASHI	31	66.0	0.034	± 0.013	0.012	± 0.011
Gifu, GIFU	31	129.0	0.022	± 0.0085	0.011	± 0.010
Shizuoka, SHIZUOKA	32	254.0	0.000	± 0.013	0.013	± 0.012
Nagoya, AICHI	31	110.2	0.046	± 0.015	0.019	± 0.015
Otsu, SHIGA	31	146.0	0.040	± 0.021	0.013	± 0.012
Tsu, MIE	31	97.0	0.026	± 0.014	0.002	± 0.011
Kyoto, KYOTO	32	145.0	0.018	± 0.0098	0.000	± 0.012
Kobe, HYOGO	31	108.3	0.0085	± 0.0074	0.020	± 0.014
Nara, NARA	31	119.3	0.013	± 0.0097	0.014	± 0.010
Wakayama, WAKAYAMA	31	106.0	0.039	± 0.014	0.020	± 0.012
Tottori, TOTTORI	31	103.2	0.070	± 0.011	0.000	± 0.011
Matsue, SHIMANE	31	108.4	0.012	± 0.0079	0.014	± 0.0082

Location	Duration (days)	Precipitation (mm)	^{90}Sr		^{137}Cs	
			(MBq/km 2)	(MBq/km 2)	(MBq/km 2)	(MBq/km 2)
Hiroshima, HIROSHIMA	31	127.5	0.099	\pm 0.016	0.025	\pm 0.015
Ishii-machi, TOKUSHIMA	31	62.2	0.030	\pm 0.018	0.37	\pm 0.031
Takamatsu, KAGAWA	31	77.0	0.012	\pm 0.0070	0.003	\pm 0.011
Matsuyama, EHIME	31	132.5	0.007	\pm 0.013	0.015	\pm 0.012
Dazaifu, FUKUOKA	31	142.4	0.007	\pm 0.011	0.008	\pm 0.011
Saga, SAGA	31	159.1	0.030	\pm 0.0087	0.007	\pm 0.011
Nagasaki, NAGASAKI	31	100.0	0.021	\pm 0.0090	0.018	\pm 0.012
Uto, KUMAMOTO	31	263.4	0.0094	\pm 0.0074	0.000	\pm 0.010
Ooita, OITA	31	180.0	0.021	\pm 0.011	0.0094	\pm 0.0098
Miyazaki, MIYAZAKI	31	121.8	0.001	\pm 0.012	0.000	\pm 0.010
Yonagusuku-machi, Okinawa	31	285.1	0.007	\pm 0.014	0.009	\pm 0.011
Aomori, AOMORI	30	40.2	0.050	\pm 0.0094	0.059	\pm 0.019
May, 1997						
Sapporo, HOKKAIDOU	33	86.5	0.086	\pm 0.022	0.19	\pm 0.031
Aomori, AOMORI	34	94.5	0.066	\pm 0.010	0.078	\pm 0.020
Morioka, IWATE	33	152.8	0.023	\pm 0.018	0.060	\pm 0.016
Onagawa-machi, MIYAGI	33	148.0	0.029	\pm 0.011	0.019	\pm 0.013
Yamagata, YAMAGATA	33	134.3	0.025	\pm 0.012	0.019	\pm 0.011
Ookuma-machi, FUKUSHIMA	33	196.2	0.031	\pm 0.014	0.006	\pm 0.011
Mito, IBARAKI	33	214.5	0.035	\pm 0.013	0.070	\pm 0.016
Kawachi-machi, TOCHIGI	33	289.7	0.027	\pm 0.0073	0.026	\pm 0.013
Urawa, SAITAMA	33	238.5	0.017	\pm 0.0070	0.038	\pm 0.011
Ichihara, CHIBA	33	185.7	0.029	\pm 0.015	0.055	\pm 0.016
Shinjuku, TOKYO	33	204.0	0.018	\pm 0.013	0.000	\pm 0.012
Yokohama, KANAGAWA	30	163.0	0.065	\pm 0.016	0.024	\pm 0.014
Kosugi-machi, TOYAMA	33	1191.1	0.0061	\pm 0.0064	0.023	\pm 0.014
Fukui, FUKUI	33	256.0	0.062	\pm 0.057	0.000	\pm 0.059

Location	Duration (days)	Precipitation (mm)	^{90}Sr		^{137}Cs	
			(MBq/km 2)	(MBq/km 2)	(MBq/km 2)	(MBq/km 2)
Koufu, YAMANASHI	33	70.0	0.031	\pm 0.013	0.012	\pm 0.011
Gifu, Gifu	33	218.5	0.0097	\pm 0.077	0.017	\pm 0.011
Shizuoka, SHIZUOKA	32	113.5	0.007	\pm 0.015	0.043	\pm 0.013
Nagoya, AICHI	33	151.8	0.002	\pm 0.011	0.030	\pm 0.016
Otsu, SHIGA	33	175.6	0.017	\pm 0.014	0.018	\pm 0.013
Tsu, MIE	33	114.0	0.000	\pm 0.011	0.026	\pm 0.014
Kyoto, KYOTO	31	159.5	0.035	\pm 0.011	0.000	\pm 0.011
Kobe, HYOUGO	31	143.8	0.030	\pm 0.0083	0.000	\pm 0.010
Nara, NARA	33	206.3	0.009	\pm 0.010	0.020	\pm 0.012
Wakayama, WAKAYAMA	32	156.5	0.010	\pm 0.013	0.013	\pm 0.012
Tottori, TOTTORI	33	211.8	0.043	\pm 0.0095	0.050	\pm 0.015
Matsue, SHIMANE	32	208.3	0.023	\pm 0.017	0.050	\pm 0.010
Hirosshima, HIROSHIMA	33	265.4	0.15	\pm 0.018	0.000	\pm 0.015
Ishii-machi, TOKUSHIMA	28	94.8	0.000	\pm 0.048	0.007	\pm 0.023
Takamatsu, KAGAWA	33	61.5	0.031	\pm 0.0083	0.000	\pm 0.010
Matsuyama, EHIME	33	120.5	0.006	\pm 0.012	0.013	\pm 0.012
Dazaifu, FUKUOKA	33	231.3	0.003	\pm 0.012	0.026	\pm 0.011
Saga, SAGA	33	312.5	0.017	\pm 0.011	0.000	\pm 0.011
Nagasaki, NAGASAKI	33	125.0	0.026	\pm 0.010	0.000	\pm 0.010
Uto, KUMAMOTO	33	146.8	0.017	\pm 0.013	0.0044	\pm 0.0094
Ooita, OITA	33	104.2	0.019	\pm 0.011	0.003	\pm 0.010
Miyazaki, MIYAZAKI	33	95.3	0.000	\pm 0.012	0.0000	\pm 0.0095
Yonagusuku-machi, Okinawa	31	268.1	0.019	\pm 0.0074	0.000	\pm 0.012
June, 1997						
Sapporo, HOKKAIDOU	30	21.5	0.017	\pm 0.0073	0.028	\pm 0.013
Aomori, AOMORI	30	113.3	0.038	\pm 0.0088	0.014	\pm 0.017
Morioka, IWATE	30	139.4	0.0000	\pm 0.020	0.000	\pm 0.011

Location	Duration (days)	Precipitation (mm)	^{90}Sr		^{137}Cs	
			(MBq/km 2)	(MBq/km 2)	(MBq/km 2)	(MBq/km 2)
Onagawa-machi, MIYAGI	29	366.5	0.0039	\pm 0.0075	0.0000	\pm 0.0093
Yamagata, YAMAGATA	31	267.5	0.012	\pm 0.0072	0.0012	\pm 0.0094
Ookuma-machi, FUKUSHIMA	30	174.4	0.018	\pm 0.011	0.030	\pm 0.011
Mito, IBARAKI	30	138.0	0.009	\pm 0.011	0.006	\pm 0.012
Kawachi-machi, TOCHIGI	30	185.1	0.0036	\pm 0.0066	0.009	\pm 0.012
Maebashi, GUNMA	30	105.5	0.015	\pm 0.0068	0.005	\pm 0.014
Urawa, SAITAMA	30	185.2	0.010	\pm 0.014	0.030	\pm 0.018
Ichihara, CHIBA	30	119.7	0.012	\pm 0.012	0.009	\pm 0.012
Shinjuku, TOKYO	30	211.4	0.020	\pm 0.014	0.010	\pm 0.016
Yokohama, KANAGAWA	32	193.6	0.017	\pm 0.013	0.017	\pm 0.011
Kosugi-machi, TOYAMA	30	155.6	0.018	\pm 0.012	0.012	\pm 0.012
Fukui, FUKUI	26	448.1	0.000	\pm 0.028	0.000	\pm 0.050
Koufu, YAMANASHI	30	142.5	0.026	\pm 0.013	0.001	\pm 0.011
Gifu, GIFU	30	92.5	0.019	\pm 0.011	0.016	\pm 0.013
Shizuoka, SHIZUOKA	30	274.5	0.017	\pm 0.0088	0.012	\pm 0.011
Nagoya, AICHI	31	142.0	0.040	\pm 0.016	0.019	\pm 0.014
Ootsu, SHIGA	30	156.2	0.0000	\pm 0.0064	0.027	\pm 0.012
Tsu, MIE	30	137.5	0.014	\pm 0.0088	0.004	\pm 0.011
Kyoto, KYOTO	29	112.0	0.037	\pm 0.010	0.000	\pm 0.011
Kobe, HYOUGO	32	53.1	0.0021	\pm 0.0078	0.000	\pm 0.012
Nara, NARA	30	128.5	0.019	\pm 0.011	0.000	\pm 0.011
Wakayama, WAKAYAMA	30	93.0	0.012	\pm 0.016	0.006	\pm 0.012
Tottori, TOTTORI	30	125.5	0.037	\pm 0.0090	0.036	\pm 0.014
Matsue, SHIMANE	30	130.0	0.013	\pm 0.0072	0.016	\pm 0.0076
Hiroshima, HIROSHIMA	29	158.9	0.17	\pm 0.040	0.000	\pm 0.015
Ishii-machi, TOKUSHIMA	35	133.1	0.038	\pm 0.014	0.029	\pm 0.039
Takamatsu, KAGAWA	30	44.5	0.0095	\pm 0.0068	0.001	\pm 0.011

Location	Duration (days)	Precipitation (mm)	^{90}Sr		^{137}Cs	
			(MBq/km ²)	(MBq/km ²)	(MBq/km ²)	(MBq/km ²)
Matsuyama, EHIME	30	132.0	0.000	\pm 0.012	0.006	\pm 0.011
Dazaifu, FUKUOKA	30	166.7	0.011	\pm 0.0083	0.016	\pm 0.011
Saga, SAGA	30	231.4	0.004	\pm 0.010	0.010	\pm 0.011
Nagasaki, NAGASAKI	30	357.5	0.013	\pm 0.0097	0.000	\pm 0.012
Uto, KUMAMOTO	30	256.3	0.013	\pm 0.0068	0.0000	\pm 0.0082
Ooita, OITA	30	200.3	0.000	\pm 0.012	0.000	\pm 0.011
Miyazaki, MIYAZAKI	30	477.6	0.013	\pm 0.014	0.017	\pm 0.011
Yonagusuku-machi, Okinawa	32	312.5	0.018	\pm 0.0096	0.000	\pm 0.019
July, 1997						
Sapporo, HOKKAIDOU	32	76.0	0.014	\pm 0.0078	0.0000	\pm 0.011
Aomori, AOMORI	31	16.4	0.011	\pm 0.0068	0.007	\pm 0.016
Morioka, IWATE	32	57.4	0.0000	\pm 0.012	0.000	\pm 0.011
Onagawa-machi, MIYAGI	32	68.5	0.0075	\pm 0.0075	0.004	\pm 0.010
Yamagata, YAMAGATA	32	139.7	0.0023	\pm 0.0073	0.016	\pm 0.012
Ookuma-machi, FUKUSHIMA	32	71.5	0.0036	\pm 0.0065	0.000	\pm 0.012
Mito, IBARAKI	32	73.0	0.020	\pm 0.012	0.0000	\pm 0.0097
Kawachi-machi, TOCHIGI	32	199.8	0.011	\pm 0.0078	0.0000	\pm 0.0091
Maebashi, GUNMA	32	178.0	0.0095	\pm 0.0070	0.005	\pm 0.011
Urawa, SAITAMA	32	128.0	0.023	\pm 0.017	0.006	\pm 0.010
Ichihara, CHIBA	32	193.2	0.021	\pm 0.014	0.009	\pm 0.014
Shinjuku, TOKYO	32	189.3	0.012	\pm 0.013	0.008	\pm 0.015
Yokohama, KANAGAWA	32	132.2	0.019	\pm 0.012	0.028	\pm 0.014
Kosugi-machi, TOYAMA	32	388.6	0.014	\pm 0.0068	0.000	\pm 0.012
Fukui, FUKUI	41	534.3	0.046	\pm 0.039	0.084	\pm 0.054
Koufu, YAMANASHI	32	175.0	0.030	\pm 0.014	0.015	\pm 0.012
Gifu, GIFU	32	615.5	0.014	\pm 0.0081	0.012	\pm 0.011
Shizuoka, SHIZUOKA	32	400.5	0.044	\pm 0.011	0.0000	\pm 0.0093

Location	Duration (days)	Precipitation (mm)	^{90}Sr		^{137}Cs	
			(MBq/km 2)	(MBq/km 2)	(MBq/km 2)	(MBq/km 2)
Nagoya, AICHI	31	436.2	0.0095	\pm 0.0063	0.022	\pm 0.012
Ootsu, SHIGA	32	387.8	0.0000	\pm 0.0062	0.007	\pm 0.011
Tsu, MIE	32	306.5	0.011	\pm 0.0095	0.019	\pm 0.012
Kyoto, KYOTO	32	334.5	0.000	\pm 0.011	0.0000	\pm 0.0096
Kobe, HYOUGO	32	454.2	0.034	\pm 0.015	0.036	\pm 0.014
Nara, NARA	32	462.5	0.0000	\pm 0.0088	0.000	\pm 0.011
Wakayama, WAKAYAMA	32	229.0	0.023	\pm 0.015	0.021	\pm 0.012
Tottori, TOTTORI	32	355.5	0.055	\pm 0.013	0.014	\pm 0.016
Matsue, SHIMANE	32	549.2	0.000	\pm 0.011	0.0043	\pm 0.0073
Hirosshima, HIROSHIMA	33	435.2	0.089	\pm 0.040	0.008	\pm 0.012
Takamatsu, KAGAWA	32	225.5	0.015	\pm 0.012	0.003	\pm 0.012
Matsuyama, EHIME	32	315.5	0.000	\pm 0.012	0.004	\pm 0.011
Dazaifu, FUKUOKA	32	593.6	0.017	\pm 0.0086	0.014	\pm 0.012
Saga, SAGA	32	635.3	0.010	\pm 0.013	0.023	\pm 0.013
Nagasaki, NAGASAKI	32	631.0	0.002	\pm 0.015	0.003	\pm 0.011
Uto, KUMAMOTO	32	959.7	0.006	\pm 0.015	0.005	\pm 0.012
Ooita, OITA	32	426.7	0.000	\pm 0.013	0.000	\pm 0.011
Miyazaki, MIYAZAKI	32	228.3	0.012	\pm 0.012	0.0000	\pm 0.0092
Yonagusuku-machi, Okinawa	36	36.0	0.012	\pm 0.0090	0.000	\pm 0.011
August, 1997						
Sapporo, HOKKAIDOU	32	186.5	0.0094	\pm 0.0086	0.0000	\pm 0.015
Aomori, AOMORI	33	143.1	0.015	\pm 0.0072	0.022	\pm 0.016
Morioka, IWATE	32	118.4	0.027	\pm 0.0095	0.018	\pm 0.015
Onagawa-machi, MIYAGI	32	13.0	0.0094	\pm 0.0072	0.015	\pm 0.011
Yamagata, YAMAGATA	32	38.3	0.000	\pm 0.015	0.000	\pm 0.011
Ookuma-machi, FUKUSHIMA	32	48.5	0.000	\pm 0.012	0.013	\pm 0.013
Mito, IBARAKI	32	48.5	0.028	\pm 0.014	0.004	\pm 0.012

Location	Duration (days)	Precipitation (mm)	^{90}Sr		^{137}Cs	
			(MBq/km ²)	(MBq/km ²)	(MBq/km ²)	(MBq/km ²)
Kawachi-machi, TOCHIGI	32	133.1	0.0000	\pm 0.0062	0.0067	\pm 0.0099
Maebashi, GUNMA	32	137.5	0.000	\pm 0.012	0.008	\pm 0.010
Urawa, SAITAMA	32	55.8	0.023	\pm 0.0099	0.008	\pm 0.014
Ichihara, CHIBA	31	121.5	0.037	\pm 0.016	0.020	\pm 0.016
Shinjuku, TOKYO	32	95.0	0.009	\pm 0.015	0.015	\pm 0.015
Yokohama, KANAGAWA	29	77.6	0.018	\pm 0.0094	0.018	\pm 0.013
Kosugi-machi, TOYAMA	32	93.6	0.015	\pm 0.0070	0.003	\pm 0.011
Fukui, FUKUI	28	110.6	0.000	\pm 0.064	0.000	\pm 0.059
Koufu, YAMANASHI	32	52.0	0.0022	\pm 0.0076	0.000	\pm 0.015
Gifu, GIFU	32	100.5	0.0000	\pm 0.0092	0.012	\pm 0.019
Shizuoka, SHIZUOKA	32	47.5	0.015	\pm 0.0064	0.000	\pm 0.010
Nagoya, AICHI	32	55.0	0.019	\pm 0.010	0.010	\pm 0.011
Ootsu, SHIGA	32	122.4	0.011	\pm 0.0085	0.008	\pm 0.012
Tsu, MIE	32	52.0	0.0056	\pm 0.0059	0.018	\pm 0.016
Kyoto, KYOTO	33	114.5	0.016	\pm 0.014	0.0036	\pm 0.0098
Kobe, HYOUGO	30	80.3	0.0090	\pm 0.0080	0.000	\pm 0.011
Nara, NARA	32	160.4	0.029	\pm 0.014	0.000	\pm 0.010
Wakayama, WAKAYAMA	32	82.0	0.020	\pm 0.020	0.015	\pm 0.012
Matsue, SHIMANE	32	207.6	0.023	\pm 0.0097	0.0000	\pm 0.0062
Hiroshima, HIROSHIMA	32	141.5	0.027	\pm 0.012	0.0076	\pm 0.0088
Takamatsu, KAGAWA	32	50.0	0.011	\pm 0.0081	0.006	\pm 0.011
Matsuyama, EHIME	32	55.5	0.0063	\pm 0.0075	0.0000	\pm 0.0080
Dazaifu, FUKUOKA	32	321.6	0.003	\pm 0.012	0.012	\pm 0.017
Saga, SAGA	32	308.8	0.0000	\pm 0.0090	0.010	\pm 0.012
Nagasaki, NAGASAKI	32	189.0	0.026	\pm 0.029	0.000	\pm 0.015
Uto, KUMAMOTO	32	167.0	0.019	\pm 0.013	0.012	\pm 0.0090
Ooita, OITA	32	156.7	0.015	\pm 0.0081	0.007	\pm 0.015

Location	Duration (days)	Precipitation (mm)	^{90}Sr		^{137}Cs	
			(MBq/km 2)	(MBq/km 2)	(MBq/km 2)	(MBq/km 2)
Miyazaki, MIYAZAKI	32	331.6	0.000	± 0.010	0.008	± 0.011
Yonagusuku-machi, Okinawa	29	454.5	0.028	± 0.0092	0.063	± 0.025
September, 1997						
Sapporo, HOKKAIDOU	31	120.0	0.0012	± 0.0060	0.0000	± 0.012
Aomori, AOMORI	31	103.6	0.025	± 0.011	0.000	± 0.013
Morioka, IWATE	31	194.8	0.0040	± 0.0063	0.019	± 0.016
Onagawa-machi, MIYAGI	31	240.0	0.016	± 0.0080	0.0000	± 0.0094
Yamagata, YAMAGATA	31	187.7	0.013	± 0.0063	0.000	± 0.012
Ookuma-machi, FUKUSHIMA	31	240.8	0.0000	± 0.0067	0.011	± 0.016
Mito, IBARAKI	31	133.0	0.0053	± 0.0083	0.000	± 0.015
Kawachi-machi, TOCHIGI	31	199.1	0.017	± 0.0074	0.000	± 0.015
Maebashi, GUNMA	31	314.0	0.002	± 0.012	0.0058	± 0.0095
Urawa, SAITAMA	31	241.4	0.016	± 0.020	0.032	± 0.012
Ichihara, CHIBA	31	162.8	0.024	± 0.014	0.000	± 0.013
Shinjuku, TOKYO	31	245.8	0.011	± 0.011	0.000	± 0.010
Yokohama, KANAGAWA	34	190.7	0.038	± 0.013	0.003	± 0.012
Kosugi-machi, TOYAMA	31	243.7	0.013	± 0.010	0.018	± 0.011
Fukui, FUKUI	30	204.7	0.057	± 0.059	0.000	± 0.046
Koufu, YAMANASHI	31	115.5	0.036	± 0.011	0.017	± 0.016
Gifu, GIFU	31	183.0	0.009	± 0.012	0.005	± 0.013
Shizuoka, SHIZUOKA	31	175.0	0.0076	± 0.0065	0.0044	± 0.0091
Nagoya, AICHI	31	189.9	0.034	± 0.015	0.011	± 0.014
Ootsu, SHIGA	31	194.3	0.021	± 0.0082	0.0091	± 0.0091
Tsu, MIE	31	189.0	0.019	± 0.0070	0.050	± 0.018
Kyoto, KYOTO	31	206.0	0.006	± 0.012	0.0000	± 0.0097
Kobe, HYOUGO	33	228.3	0.006	± 0.011	0.000	± 0.011

Location	Duration (days)	Precipitation (mm)	^{90}Sr		^{137}Cs	
			(MBq/km 2)	(MBq/km 2)	(MBq/km 2)	(MBq/km 2)
Nara, NARA	31	216.2	0.015	± 0.013	0.11	± 0.020
Wakayama, WAKAYAMA	31	240.5	0.000	± 0.017	0.005	± 0.013
Tottori, TOTTORI	31	289.1	0.056	± 0.023	0.043	± 0.020
Matsue, SHIMANE	31	352.7	0.0080	± 0.0086	0.014	± 0.0078
Hirosshima, HIROSHIMA	30	237.2	0.027	± 0.0080	0.0000	± 0.0097
Takamatsu, KAGAWA	31	162.0	0.0022	± 0.0078	0.002	± 0.010
Matsuyama, EHIME	31	213.0	0.021	± 0.0084	0.034	± 0.012
Dazaifu, FUKUOKA	31	211.8	0.000	± 0.010	0.013	± 0.017
Saga, SAGA	31	270.1	0.012	± 0.012	0.013	± 0.012
Nagasaki, NAGASAKI	31	271.0	0.019	± 0.014	0.0024	± 0.0099
Uto, KUMAMOTO	31	71.8	0.021	± 0.0073	0.008	± 0.016
Ooita, OITA	31	323.1	0.0040	± 0.0069	0.011	± 0.017
Miyazaki, MIYAZAKI	31	473.8	0.008	± 0.011	0.0000	± 0.0097
Yonagusuku-machi, Okinawa	30	103.5	0.065	± 0.0084	0.000	± 0.058

6. Results

(1)-2 Strontium-90 and Cesium-137 in Rain and Dry Fallout(for WHO program)
(from Apr. 1997 to Sep. 1997)

-continued from No. 120 of this publication-

Table (1)-2 : Strontium-90 and Cesium-137 in Rain and Dry Fallout

Location	Duration (days)	Precipitation (mm)	^{90}Sr		^{137}Cs	
			(MBq/km ²)	(MBq/km ²)	(MBq/km ²)	(MBq/km ²)
April, 1997						
Akita, AKITA	31	98.6	0.044	\pm 0.016	0.018	\pm 0.013
Chiba, CHIBA	32	105.3	0.022	\pm 0.020	0.020	\pm 0.011
Niigata, NIIGATA	31	132.4	0.052	\pm 0.0091	0.034	\pm 0.014
Kanazawa, ISHIKAWA	31	88.5	0.016	\pm 0.011	0.043	\pm 0.015
Nagano, NAGANO	32	60.7	0.032	\pm 0.0091	0.000	\pm 0.011
Osaka, OSAKA	32	128.6	0.038	\pm 0.012	0.023	\pm 0.013
Okayama, OKAYAMA	31	88.0	0.0078	\pm 0.0078	0.025	\pm 0.012
Yamaguchi, YAMAGUCHI	31	148.0	0.022	\pm 0.0086	0.016	\pm 0.012
Kochi, KOCHI	31	132.1	0.055	\pm 0.011	0.015	\pm 0.012
Kagoshima, KAGOSHIMA	31	95.5	0.034	\pm 0.0086	0.050	\pm 0.015
May, 1997						
Akita, AKITA	33	177.4	0.036	\pm 0.014	0.047	\pm 0.015
Chiba, CHIBA	32	175.3	0.056	\pm 0.025	0.044	\pm 0.013
Niigata, NIIGATA	33	182.9	0.043	\pm 0.0087	0.036	\pm 0.014
Kanazawa, ISHIKAWA	31	129.5	0.046	\pm 0.014	0.060	\pm 0.015
Nagano, NAGANO	32	67.9	0.035	\pm 0.013	0.000	\pm 0.015
Osaka, OSAKA	33	102.6	0.042	\pm 0.023	0.018	\pm 0.013
Okayama, OKAYAMA	33	120.6	0.0022	\pm 0.0072	0.030	\pm 0.011
Yamaguchi, YAMAGUCHI	32	357.0	0.028	\pm 0.0083	0.011	\pm 0.012
Kochi, KOCHI	33	142.4	0.071	\pm 0.017	0.019	\pm 0.012
Kagoshima, KAGOSHIMA	31	116.0	0.012	\pm 0.0078	0.024	\pm 0.012
June, 1997						
Akita, AKITA	30	161.7	0.023	\pm 0.014	0.016	\pm 0.011
Chiba, CHIBA	30	109.0	0.016	\pm 0.012	0.032	\pm 0.012
Niigata, NIIGATA	30	256.2	0.027	\pm 0.0077	0.016	\pm 0.012
Kanazawa, ISHIKAWA	32	167.0	0.011	\pm 0.010	0.000	\pm 0.010

Location	Duration (days)	Precipitation (mm)	^{90}Sr		^{137}Cs	
			(MBq/km 2)	(MBq/km 2)	(MBq/km 2)	(MBq/km 2)
Nagano, NAGANO	31	75.6	0.010	± 0.020	0.000	± 0.026
Osaka, OSAKA	30	62.6	0.007	± 0.014	0.005	± 0.011
Okayama, OKAYAMA	30	100.1	0.018	± 0.0085	0.0079	± 0.0082
Yamaguchi, YAMAGUCHI	31	210.0	0.023	± 0.0082	0.004	± 0.012
Kochi, KOCHI	30	313.2	0.071	± 0.011	0.015	± 0.014
Kagoshima, KAGOSHIMA	29	306.0	0.0098	± 0.0073	0.006	± 0.011
July, 1997						
Akita, AKITA	32	118.5	0.026	± 0.0097	0.005	± 0.011
Chiba, CHIBA	32	155.5	0.045	± 0.029	0.016	± 0.011
Niigata, NIIGATA	32	203.9	0.000	± 0.012	0.000	± 0.014
Kanazawa, ISHIKAWA	32	423.0	0.026	± 0.0076	0.001	± 0.011
Nagano, NAGANO	31	116.2	0.015	± 0.011	0.004	± 0.015
Osaka, OSAKA	32	276.6	0.041	± 0.043	0.005	± 0.013
Okayama, OKAYAMA	32	256.0	0.000	± 0.012	0.000	± 0.011
Yamaguchi, YAMAGUCHI	32	464.0	0.0006	± 0.0081	0.000	± 0.011
Kochi, KOCHI	32	242.5	0.062	± 0.011	0.026	± 0.012
Kagoshima, KAGOSHIMA	35	371.0	0.031	± 0.0088	0.021	± 0.013
August, 1997						
Akita, AKITA	32	123.2	0.036	± 0.017	0.011	± 0.012
Chiba, CHIBA	32	26.6	0.013	± 0.016	0.005	± 0.010
Niigata, NIIGATA	32	100.0	0.013	± 0.0015	0.000	± 0.014
Kanazawa, ISHIKAWA	30	105.5	0.021	± 0.0076	0.014	± 0.011
Nagano, NAGANO	32	83.9	0.029	± 0.014	0.001	± 0.013
Osaka, OSAKA	32	119.2	0.010	± 0.0091	0.000	± 0.011
Okayama, OKAYAMA	32	32.1	0.012	± 0.015	0.004	± 0.010
Yamaguchi, YAMAGUCHI	32	235.0	0.014	± 0.013	0.000	± 0.011
Kochi, KOCHI	32	285.7	0.70	± 0.011	0.000	± 0.011

Location	Duration (days)	Precipitation (mm)	^{90}Sr		^{137}Cs	
			(MBq/km ²)	(MBq/km ²)	(MBq/km ²)	(MBq/km ²)
Kagoshima, KAGOSHIMA	30	104.0	0.030	\pm 0.0096	0.000	\pm 0.012
September, 1997						
Akita, AKITA	31	226.2	0.020	\pm 0.0085	0.000	\pm 0.011
Chiba, CHIBA	31	131.6	0.004	\pm 0.013	0.005	\pm 0.010
Niigata, NIIGATA	31	213.5	0.039	\pm 0.017	0.017	\pm 0.016
Kanazawa, ISHIKAWA	33	244.0	0.0095	\pm 0.0066	0.000	\pm 0.011
Nagano, NAGANO	32	136.7	0.040	\pm 0.017	0.000	\pm 0.011
Osaka, OSAKA	31	201.6	0.020	\pm 0.022	0.005	\pm 0.011
Okayama, OKAYAMA	31	166.7	0.0068	\pm 0.0062	0.000	\pm 0.012
Yamaguchi, YAMAGUCHI	31	245.0	0.000	\pm 0.012	0.008	\pm 0.017
Kochi, KOCHI	31	561.9	0.030	\pm 0.014	0.003	\pm 0.012
Kagoshima, KAGOSHIMA	33	289.0	0.019	\pm 0.012	0.008	\pm 0.017

(2) Strontium-90 and Cesium-137 in Airborn Dust
 (from Apr. 1997 to Sep. 1997)

-continued from No. 120 of this publication-

Table (2) :Strontium-90 and Cesium-137 in Airborn Dust

Location	Sampling period	Absorption volume (m ³)	⁹⁰ Sr (mBq/m ³)	¹³⁷ Cs (mBq/m ³)
April-June 1997				
Morioka, IWATE	4~ 6	12,390.0	0.00056	± 0.00049 0.00031 ± 0.00029
Akita, AKITA	4~ 6	10,800.0	0.00034	± 0.00033 0.0012 ± 0.00042
Yamagata, YAMAGATA	4~ 6	12,960.0	0.00000	± 0.00024 0.00038 ± 0.00033
Ookuma-machi, FUKUSHIMA	4~ 6	10,722.3	0.00000	± 0.00029 0.00000 ± 0.00038
Mito, IBARAKI	4~ 6	10,972.0	0.00045	± 0.00034 0.00000 ± 0.00027
Kawachi-machi, TOCHIGI	4~ 5	15,010.0	0.00019	± 0.00022 0.00040 ± 0.00031
Maebashi, GUNMA	4~ 6	12,279.0	0.00006	± 0.00041 0.00000 ± 0.00030
Ichihara, CHIBA	4~ 6	10,358.0	0.00088	± 0.00057 0.00000 ± 0.00028
Niigata, NIIGATA	4~ 6	10,335.0	0.00025	± 0.00052 0.00008 ± 0.00032
Kosugi-machi, TOYAMA	4~ 6	18,557.0	0.00008	± 0.00025 0.00033 ± 0.00027
Fukui, FUKUI	4~ 6	13,078.2	0.00031	± 0.00027 0.00000 ± 0.00029
Koufu, YAMANASHI	4~ 6	11,609.0	0.00034	± 0.00044 0.00000 ± 0.00028
Nagano, NAGANO	4~ 6	11,322.0	0.00000	± 0.00041 0.00007 ± 0.00035
Gifu, GIFU	4~ 6	11,257.0	0.00070	± 0.00036 0.00020 ± 0.00031
Hamaoka-machi, SHIZUOKA	4~ 6	10,430.0	0.0010	± 0.00054 0.00000 ± 0.00038
Nagoya, AICHI	4~ 6	10,436.0	0.00014	± 0.00065 0.00007 ± 0.00036
Otsu, SHIGA	4~ 6	10,725.0	0.00000	± 0.00050 0.00000 ± 0.00033
Tsu, MIE	4~ 6	13,520.0	0.00000	± 0.00035 0.00075 ± 0.00036
Kyoto, KYOTO	4~ 6	10,287.0	0.0013	± 0.00036 0.00004 ± 0.00036
Osaka, OSAKA	4~ 6	14,230.0	0.00027	± 0.00024 0.00000 ± 0.00024
Kobe, HYOUGO	4~ 6	10,357.0	0.00013	± 0.00031 0.00066 ± 0.00042
Nara, NARA	4~ 6	10,749.7	0.00012	± 0.00051 0.00000 ± 0.00037
Wakayama, WAKAYAMA	4~ 6	10,368.0	0.00033	± 0.00057 0.00000 ± 0.00040
Tottori, TOTTORI	4~ 6	15,068.0	0.00091	± 0.00043 0.00000 ± 0.00029

Location	Sampling period	Absorption volume (m ³)	⁹⁰ Sr		¹³⁷ Cs	
			(mBq/m ³)	(mBq/m ³)	(mBq/m ³)	(mBq/m ³)
Okayama, OKAYAMA	4~ 6	12,615.0	0.00000	± 0.00047	0.00006	± 0.00030
Hiroshima, HIROSHIMA	4~ 6	10,587.0	0.00077	± 0.00055	0.00000	± 0.00035
Yamaguchi, YAMAGUCHI	4~ 6	22,091.0	0.00026	± 0.00016	0.00012	± 0.00023
Tokushima, TOKUSHIMA	4~ 6	8,400.0	0.0019	± 0.00080	0.00000	± 0.00041
Takamatsu, KAGAWA	4~ 6	15,604.1	0.00060	± 0.00039	0.00000	± 0.00026
Saga, SAGA	4~ 6	9,616.0	0.00025	± 0.00035	0.00000	± 0.00044
Nagasaki, NAGASAKI	4~ 6	10,368.0	0.00000	± 0.00051	0.00004	± 0.00037
Uto, KUMAMOTO	4~ 6	9,797.0	0.00049	± 0.00063	0.00066	± 0.00045
Ooita, OITA	4~ 6	10,364.0	0.00024	± 0.00054	0.00000	± 0.00041
Miyazaki, MIYAZAKI	4~ 6	13,795.0	0.00070	± 0.00046	0.00033	± 0.00026
June~July 1997						
Yokohama, KANAGAWA	6~ 7	7,201.0	0.00000	± 0.00038	0.00000	± 0.00057
July~September 1997						
Morioka, IWATE	7~ 9	10,294.0	0.00000	± 0.00059	0.00020	± 0.00042
Akita, AKITA	7~ 9	10,800.0	0.00000	± 0.00051	0.00051	± 0.00043
Yamagata, YAMAGATA	7~ 9	12,960.0	0.00000	± 0.00037	0.00018	± 0.00033
Ookuma-machi, FUKUSHIMA	7~ 9	10,456.0	0.00049	± 0.00058	0.00000	± 0.00034
Mito, IBARAKI	7~ 9	11,689.0	0.0012	± 0.00055	0.00000	± 0.00025
Maebashi, GUNMA	7~ 9	11,951.0	0.00000	± 0.00039	0.00000	± 0.00035
Ichihara, CHIBA	7~ 9	10,194.0	0.00007	± 0.00046	0.00016	± 0.00037
Yokohama, KANAGAWA	7~ 9	10,037.0	0.00035	± 0.00052	0.00032	± 0.00040
Niigata, NIIGATA	7~ 9	10,295.0	0.00013	± 0.00053	0.00046	± 0.00038
Kosugi-machi, TOYAMA	7~ 9	18,319.0	0.00072	± 0.00034	0.00018	± 0.00021
Fukui, FUKUI	7~ 9	12,933.0	0.00062	± 0.00044	0.00004	± 0.00034
Koufu, YAMANASHI	7~ 9	11,814.0	0.00014	± 0.00058	0.00000	± 0.00032

Location	Sampling	Absorption volume (m ³)	⁹⁰ Sr		¹³⁷ Cs	
	period		(mBq/m ³)	(mBq/m ³)	(mBq/m ³)	(mBq/m ³)
Nagano, NAGANO	7~ 9	13,002.0	0.00000	± 0.00036	0.00000	± 0.00030
Gifu, Gifu	7~ 9	11,568.0	0.00000	± 0.00048	0.00000	± 0.00032
Hameoka-machi, SHIZUOKA	7~ 9	10,060.0	0.00006	± 0.00047	0.00000	± 0.00046
Nagoya, AICHI	7~ 9	9,861.0	0.00028	± 0.00062	0.00079	± 0.00044
Gotsu, SHIGA	7~ 9	10,230.0	0.0014	± 0.00664	0.00000	± 0.00034
Tsu, MIE	7~ 9	13,460.0	0.00000	± 0.00039	0.00023	± 0.00031
Kyoto, KYOTO	7~ 9	10,365.0	0.00020	± 0.00027	0.00007	± 0.00038
Osaka, OSAKA	7~ 9	16,221.0	0.00041	± 0.00034	0.00002	± 0.00025
Kobe, HYOGO	7~ 9	10,311.0	0.00000	± 0.00057	0.00000	± 0.00033
Nara, NARA	7~ 9	10,840.7	0.00053	± 0.00055	0.00028	± 0.00040
Wakayama, WAKAYAMA	7~ 9	10,368.0	0.00033	± 0.00057	0.00000	± 0.00036
Tottori, TOTTORI	7~ 9	15,034.0	0.00059	± 0.00040	0.00005	± 0.00024
Okayama, OKAYAMA	7~ 9	15,524.0	0.00018	± 0.00040	0.00000	± 0.00024
Hirosima, HIROSHIMA	7~ 9	13,226.0	0.00042	± 0.00049	0.00010	± 0.00032
Yamaguchi, YAMAGUCHI	7~ 9	22,202.0	0.00035	± 0.00026	0.00000	± 0.00016
Tokushima, TOKUSHIMA	7~ 9	10,080.0	0.00019	± 0.00051	0.00000	± 0.00035
Takamatsu, KAGAWA	7~ 9	15,825.3	0.00047	± 0.00035	0.00000	± 0.00024
Saga, SAGA	7~ 9	10,216.6	0.00049	± 0.00030	0.00000	± 0.00032
Nagasaki, NAGASAKI	7~ 9	10,368.0	0.00012	± 0.00072	0.00000	± 0.00031
Uto, KUMAMOTO	7~ 9	11,191.0	0.00069	± 0.00051	0.00038	± 0.00058
Ooita, OITA	7~ 9	10,362.0	0.00023	± 0.00027	0.00008	± 0.00040
Miyazaki, MIYAZAKI	7~ 9	13,089.0	0.0010	± 0.00049	0.00000	± 0.00030
August, 1997						
Kawachi-machi, TOCHIGI	8~ 9	15,302.0	0.00060	± 0.00039	0.00032	± 0.00026

(3) Strontium-90 and cesium-137 in Service Water
 (from Apr. 1997 to Sep. 1997)

(20)

-continued from No. 120 of this publication-

Table (3) :Strontium-90 and cesium-137 in Service Water

Location	pH	^{90}Sr		^{137}Cs		
		(mBq/ℓ)	(mBq/ℓ)	(mBq/ℓ)	(mBq/ℓ)	
(Source Water)						
June, 1996						
Urawa, SAITAMA	7.6	0.00	± 0.23	0.000	± 0.062	
Kisarazu, CHIBA	7.4	2.0	± 0.16	0.084	± 0.077	
Katsushika, TOKYO	7.1	1.6	± 0.25	0.31	± 0.095	
Tsukui-machi, KANAGAWA	7.4	0.48	± 0.060	0.082	± 0.060	
Nagano, NAGANO	7.24	1.0	± 0.14	0.095	± 0.060	
Inuyama, AICHI	7.1	2.0	± 0.11	0.13	± 0.064	
Moriguchi, OSAKA	7.0	2.8	± 0.13	0.000	± 0.056	
Fukuoka, FUKUOKA	9.40	1.5	± 0.14	0.000	± 0.053	
July, 1996						
Sapporo, HOKKAIDOU	7.4	1.5	± 0.10	0.23	± 0.075	
August, 1996						
Kyoto, KYOTO	8.35	3.0	± 0.22	0.11	± 0.064	
(Tap Water)						
June, 1996						
Wakkanai, HOKKAIDOU	6.9	1.1	± 0.09	0.000	± 0.049	
Aomori, AOMORI	7.4	1.3	± 0.10	0.34	± 0.084	
Morioka, IWATE	7.1	0.95	± 0.082	0.000	± 0.047	
Yamagata, YAMAGATA	7.1	1.7	± 0.17	0.054	± 0.065	
Fukushima, FUKUSHIMA	8.03	2.9	± 0.20	0.086	± 0.055	
Mito, IBARAKI	7.5	1.6	± 0.15	0.018	± 0.056	
Maebashi, GUNMA	7.1	1.2	± 0.13	0.19	± 0.081	
Urawa, SAITAMA	6.9	1.4	± 0.11	0.11	± 0.079	
Ichihara, CHIBA	7.8	2.1	± 0.17	0.000	± 0.076	
Katsushika, TOKYO	7.1	1.8	± 0.21	0.12	± 0.079	

Location	pH	^{90}Sr		^{137}Cs	
		(mBq/ ℓ)	(mBq/ ℓ)	(mBq/ ℓ)	(mBq/ ℓ)
Yokohama, KANAGAWA	7.4	0.56	\pm 0.063	0.052	\pm 0.062
Niigata, NIIGATA	7.19	2.4	\pm 0.13	0.25	\pm 0.087
Kosugi-machi, TOYAMA	6.4	0.25	\pm 0.082	0.13	\pm 0.061
Kanazawa, ISHIKAWA	6.6	2.0	\pm 0.17	0.012	\pm 0.065
Fukui, FUKUI	6.30	0.46	\pm 0.096	0.000	\pm 0.053
Koufu, YAMANASHI	7.2	1.4	\pm 0.14	0.012	\pm 0.067
Nagano, NAGANO	7.42	0.70	\pm 0.12	0.012	\pm 0.051
Gifu, GIFU	7.03	1.6	\pm 0.18	0.000	\pm 0.045
Shizuoka, SHIZUOKA	7.4	0.78	\pm 0.086	0.041	\pm 0.074
Nagoya, AICHI	6.6	2.1	\pm 0.12	0.086	\pm 0.061
Ootsu, SHIGA	6.7	3.4	\pm 0.16	0.073	\pm 0.061
Tsu, MIE	7.1	2.6	\pm 0.12	0.058	\pm 0.068
Osaka, OSAKA	7.2	3.2	\pm 0.15	0.000	\pm 0.054
Kobe, HYOUGO	7.21	1.1	\pm 0.15	0.060	\pm 0.049
Nara, NARA	7.1	2.8	\pm 0.14	0.025	\pm 0.053
Tottori, TOTTORI	6.9	3.2	\pm 0.20	0.012	\pm 0.055
Matsue, SHIMANE	—	3.6	\pm 0.23	0.041	\pm 0.074
Okayama, OKAYAMA	6.9	2.5	\pm 0.19	0.060	\pm 0.076
Hiroshima, HIROSHIMA	7.2	2.2	\pm 0.15	0.053	\pm 0.062
Ube, YAMAGUCHI	6.0	2.6	\pm 0.29	0.030	\pm 0.054
Takamatsu, KAGAWA	7.60	2.4	\pm 0.12	0.14	\pm 0.058
Matsuyama, EHIME	7.7	1.5	\pm 0.09	0.078	\pm 0.062
Kochi, KOCHI	7.6	1.7	\pm 0.11	0.000	\pm 0.054
Fukuoka, FUKUOKA	6.95	2.6	\pm 0.20	0.000	\pm 0.058
Saga, SAGA	7.58	1.8	\pm 0.13	0.000	\pm 0.043
Nagasaki, NAGASAKI	6.9	1.0	\pm 0.08	0.000	\pm 0.051

Location	pH	^{90}Sr		^{137}Cs	
		(mBq/m ³)	(mBq/m ³)	(mBq/m ³)	(mBq/m ³)
Uto, KUMAMOTO	7.53	0.049	\pm 0.032	0.030	\pm 0.074
Ooita, OITA	7.81	1.0	\pm 0.13	0.22	\pm 0.076
Miyazaki, MIYAZAKI	6.86	1.3	\pm 0.17	0.052	\pm 0.053
Kagoshima, KAGOSHIMA	7.7	0.65	\pm 0.11	0.21	\pm 0.084
July, 1996					
Sendai, MIYAGI	—	1.4	\pm 0.16	0.24	\pm 0.083
Akita, AKITA	6.41	2.8	\pm 0.15	0.13	\pm 0.075
Kawachi-machi, TOCHIGI	7.32	0.28	\pm 0.051	0.005	\pm 0.054
Naha, Okinawa	7.10	3.4	\pm 0.15	0.000	\pm 0.070
August, 1996					
Kyoto, KYOTO	7.56	2.8	\pm 0.23	0.006	\pm 0.059
September, 1996					
Shinguu, WAKAYAMA	6.5	1.2	\pm 0.08	0.069	\pm 0.072

(4) Strontium-90 and cesium-137 in Freshwater
 (from Apr. 1997 to Sep. 1997)

-continued from No. 120 of this publication-

Table (4) :Strontium-90 and cesium-137 in Freshwater

Location	pH	⁹⁰ Sr (mBq/ ℓ)	¹³⁷ Cs (mBq/ ℓ)
(Source Water)			
June, 1997			
Urawa, SAITAMA	7. 6	0.37	± 0.12
Kisarazu, CHIBA	7. 4	2.5	± 0.064
Katsushika, TOKYO	7. 3	2.0	± 0.063
Tsukui-machi, KANAGAWA	8. 1	0.44	± 0.051
Nagano, NAGANO	7.34	1.1	± 0.058
Inuyama, AICHI	7. 1	1.5	± 0.082
Moriguchi, OSAKA	7. 2	2.8	± 0.056
Fukuoka, FUKUOKA	7.18	2.0	± 0.054
July, 1997			
Sapporo, HOKKAIDOU	7. 1	1.7	± 0.081
August, 1997			
Kyoto, KYOTO	7. 80	2.8	± 0.049

(5) Strontium-90 and Cesium-137 in Soil

(24)

(from APR. 1997 to Sep. 1997)

-continued from No. 120 of this publication-

Table (5) Strontium-90 and Cesium-137 in Soil

Location	Sampling Depth (cm)	⁹⁰ Sr				¹³⁷ Cs			
		(Bq/kg) (dried Soil)	(MBq/km ²)		(Bq/kg) (dried Soil)	(MBq/km ²)			
May, 1997									
Tokai-mura, IBARAKI	0~ 5	9.6 ± 0.37	450	± 17	54	± 0.8	2500	± 40	
	5~20	9.5 ± 0.37	1400	± 50	16	± 0.4	2500	± 70	
Akabane-machi, AICHI	0~ 5	1.3 ± 0.09	52	± 3.7	12	± 0.4	490	± 15	
	5~20	1.1 ± 0.09	150	± 12	17	± 0.5	2300	± 60	
June, 1997									
Fukushima, FUKUSHIMA	0~ 5	5.1 ± 0.28	150	± 8	34	± 0.6	980	± 18	
	5~20	5.5 ± 0.29	510	± 27	15	± 0.4	1400	± 40	
Katsushika, TOKYO	0~ 5	0.69 ± 0.11	26	± 4.1	3.6	± 0.22	130	± 8	
	5~20	0.89 ± 0.12	88	± 12	2.6	± 0.20	260	± 19	
July, 1997									
Aomori, AOMORI	0~ 5	2.3 ± 0.12	77	± 4.0	7.6	± 0.30	260	± 10	
	5~20	2.0 ± 0.11	210	± 12	4.5	± 0.23	460	± 23	
Mutsu, AOMORI	0~ 5	3.7 ± 0.24	130	± 8	20	± 0.5	710	± 17	
	5~20	3.6 ± 0.24	320	± 21	5.5	± 0.25	480	± 22	
Yamagata, YAMAGATA	0~ 5	3.4 ± 0.15	140	± 6	20	± 0.5	850	± 20	
	5~20	1.2 ± 0.10	130	± 11	3.9	± 0.21	420	± 23	
Maebashi, GUNMA	0~ 5	0.14 ± 0.049	7.2	± 2.6	0.54	± 0.10	29	± 5.3	
	5~20	0.38 ± 0.074	69	± 13	0.49	± 0.096	89	± 17	
Ichihara, CHIBA	0~ 5	0.078 ± 0.036	3.4	± 1.6	1.2	± 0.13	55	± 56.	
	5~20	0.18 ± 0.041	30	± 6.8	2.0	± 0.16	340	± 27	
Kashiwazaki, NIIGATA	0~ 5	0.55 ± 0.10	37	± 6.9	5.5	± 0.27	360	± 18	
	5~20	0.39 ± 0.056	83	± 12	8.1	± 0.32	1700	± 70	
Nagano, NAGANO	0~ 5	2.8 ± 0.13	78	± 3.8	31	± 0.6	890	± 17	
	5~20	1.9 ± 0.12	160	± 10	7.7	± 0.30	640	± 25	
Gifu, GIFU	0~ 5	0.84 ± 0.074	24	± 2.1	11	± 0.4	310	± 10	
	5~20	0.78 ± 0.071	77	± 7.0	9.7	± 0.33	950	± 33	

Location	Sampling Depth (cm)	⁹⁰ Sr			¹³⁷ Cs		
		(Bq/kg) (dried Soil)	(MBq/km ²)	(Bq/kg) (dried Soil)	(MBq/km ²)		
Tsu, MIE	0~ 5	0.093 ± 0.038	6.8 ± 2.8	1.0 ± 0.12	75 ± 9.0		
	5~20	0.29 ± 0.050	55 ± 9.5	1.1 ± 0.13	210 ± 24		
Kyoto, KYOTO	0~ 5	1.6 ± 0.16	41 ± 4.1	3.0 ± 0.19	77 ± 5.0		
	5~20	0.93 ± 0.13	120 ± 17	5.9 ± 0.27	780 ± 36		
Osaka, OSAKA	0~ 5	0.27 ± 0.808	14 ± 4.2	2.3 ± 0.17	120 ± 9		
	5~20	0.37 ± 0.090	62 ± 15	1.1 ± 0.15	180 ± 25		
Kasai, HYOUGO	0~ 5	5.8 ± 0.19	210 ± 7	37 ± 0.7	1300 ± 20		
	5~20	0.83 ± 0.076	93 ± 8.5	4.4 ± 0.23	500 ± 26		
Kashihara, NARA	0~ 5	0.83 ± 0.076	67 ± 6.1	3.8 ± 0.22	300 ± 18		
	5~20	0.81 ± 0.11	83 ± 12	4.3 ± 0.23	440 ± 23		
Kokufu-machi, TOTTORI	0~ 5	0.19 ± 0.075	12 ± 4.7	1.3 ± 0.14	85 ± 8.8		
	5~20	0.14 ± 0.070	12 ± 5.7	0.61 ± 0.098	49 ± 7.9		
Oota, SHIMANE	0~ 5	10 ± 0.4	220 ± 8	61 ± 0.8	1300 ± 20		
	5~20	5.5 ± 0.29	270 ± 14	61 ± 0.8	3000 ± 40		
Hiroshima, HIROSHIMA	0~ 5	0.69 ± 0.11	38 ± 6.1	1.7 ± 0.14	92 ± 7.9		
	5~20	1.7 ± 0.10	290 ± 17	10 ± 0.3	1700 ± 60		
Sakaide, KAGAWA	0~ 5	2.6 ± 0.13	87 ± 4.3	14 ± 0.4	460 ± 14		
	5~20	3.1 ± 0.14	260 ± 12	4.4 ± 0.23	370 ± 19		
Matsuyama, EHIME	0~ 5	3.0 ± 0.13	50 ± 2.2	21 ± 0.5	350 ± 8		
	5~20	0.62 ± 0.069	45 ± 5.1	9.8 ± 0.34	720 ± 25		
Kochi, KOCHI	0~ 5	3.5 ± 0.22	130 ± 8	15 ± 0.4	560 ± 16		
	5~20	5.1 ± 0.27	410 ± 22	12 ± 0.4	1000 ± 30		
Fukuoka, FUKUOKA	0~ 5	4.2 ± 0.25	250 ± 14	5.1 ± 0.25	300 ± 15		
	5~20	3.0 ± 0.21	430 ± 31	4.5 ± 0.23	640 ± 33		
Obama-machi, NAGASAKI	0~ 5	2.7 ± 0.21	61 ± 4.6	54 ± 0.8	1200 ± 20		
	5~20	3.8 ± 0.24	290 ± 18	8.6 ± 0.32	670 ± 25		
Nishihara-mura, KUMAMOTO	0~ 5	4.3 ± 0.25	120 ± 7	55 ± 0.8	1500 ± 20		

Location	Sampling Depth (cm)	⁹⁰ Sr				¹³⁷ Cs			
		(Bq/kg) (dried Soil)		(MBq/km ²)		(Bq/kg) (dried Soil)		(MBq/km ²)	
		5~20	0~ 5	5~20	0~ 5	5~20	0~ 5	5~20	0~ 5
Kujuu-machi, OITA	5~20	4.3	± 0.25	240	± 14	11	± 0.4	620	± 20
	0~ 5	2.5	± 0.20	44	± 3.5	79	± 1.0	1400	± 20
	5~20	1.8	± 0.18	90	± 8.6	16	± 0.4	770	± 21
August, 1997									
Sapporo, HOKKAIDOU	0~ 5	9.8	± 0.41	360	± 15	22	± 0.5	820	± 19
	5~20	6.5	± 0.34	1200	± 60	8.2	± 0.32	1500	± 60
Takizawa-mura, IWATE	0~ 5	2.8	± 0.21	120	± 9	12	± 0.4	520	± 16
	5~20	5.1	± 0.29	700	± 40	11	± 0.4	1500	± 50
Imaichi, TOCHIGI	0~ 5	12	± 0.3	220	± 5	49	± 0.7	880	± 13
	5~20	5.0	± 0.19	110	± 4	7.2	± 0.29	160	± 7
Urawa, SAITAMA	0~ 5	1.4	± 0.09	31	± 2.0	10	± 0.3	220	± 8
	5~20	0.95	± 0.080	78	± 6.5	1.3	± 0.13	100	± 11
Yokohama, KANAGAWA	0~ 5	7.7	± 0.21	240	± 6	25	± 0.5	770	± 17
	5~20	8.6	± 0.43	1000	± 50	13	± 0.4	1500	± 50
Kosugi-machi, TOYAMA	0~ 5	0.34	± 0.12	23	± 8.1	0.73	± 0.10	50	± 7.1
	5~20	0.19	± 0.080	36	± 15	0.41	± 0.11	76	± 21
Kanazawa, ISHIKAWA	0~ 5	6.4	± 0.30	280	± 13	18	± 0.5	790	± 20
	5~20	5.9	± 0.29	910	± 45	19	± 0.5	2900	± 70
Fukui, FUKUI	0~ 5	0.36	± 0.086	32	± 7.5	3.9	± 0.23	340	± 20
	5~20	0.60	± 0.11	110	± 19	2.6	± 0.19	480	± 35
Takane-machi, YAMANASHI	0~ 5	8.4	± 0.43	220	± 11	29	± 0.6	760	± 16
	5~20	6.8	± 0.33	600	± 30	16	± 0.4	1400	± 40
Gotenba, SHIZUOKA	0~ 5	0.75	± 0.14	25	± 4.6	8.1	± 0.32	270	± 11
	5~20	0.19	± 0.096	17	± 8.6	2.1	± 0.16	190	± 15
Yasu-machi, FUKUOKA	0~ 5	0.15	± 0.045	7.5	± 2.2	0.51	± 0.091	25	± 4.5

Location	Sampling Depth (cm)	⁹⁰ Sr				¹³⁷ Cs			
		(Bq/kg) (dried Soil)	(MBq/km ²)		(Bq/kg) (dried Soil)	(MBq/km ²)			
Asahi-machi, OKAYAMA	5~20	0.10 ± 0.042	15	± 6.2	0.41 ± 0.082	59	± 12		
	0~ 5	1.3 ± 0.15	59	± 6.6	1.5 ± 0.14	66	± 6.3		
Hagi, YAMAGUCHI	5~20	0.77 ± 0.12	79	± 12	0.57 ± 0.096	58	± 9.8		
	0~ 5	1.0 ± 0.08	66	± 5.2	6.5 ± 0.28	400	± 17		
Sadohara-machi, MIYAZAKI	5~20	1.5 ± 0.10	330	± 22	5.4 ± 0.25	1200	± 60		
	0~ 5	0.90 ± 0.12	64	± 8.3	6.7 ± 0.28	470	± 20		
Naha, Okinawa	5~20	1.0 ± 0.13	100	± 13	6.9 ± 0.28	690	± 29		
	0~ 5	0.87 ± 0.12	59	± 7.9	3.8 ± 0.22	260	± 15		
September, 1997		5~20	1.2 ± 0.14	220	± 28	2.3 ± 0.17	450	± 33	
Iwadeyama-machi, MIYAGI	0~ 5	1.6 ± 0.16	49	± 4.9	5.0 ± 0.26	150	± 8		
	5~20	1.4 ± 0.15	200	± 22	1.8 ± 0.17	260	± 25		
Saga, SAGA	0~ 5	0.48 ± 0.10	15	± 3.2	4.3 ± 0.24	140	± 8		
	5~20	0.70 ± 0.12	130	± 23	3.8 ± 0.23	720	± 44		
Kaimon-machi, KAGOSHIMA	0~ 5	0.22 ± 0.072	12	± 4.1	0.48 ± 0.11	27	± 6.1		
	5~20	0.30 ± 0.081	39	± 10	0.95 ± 0.14	120	± 18		

(6) Strontium-90 and Cesium-137 in Sea Water

(from Apr. 1997 to Sep. 1997)

-continued from No. 120 of this publication-

Table (6) Strontium-90 and Cesium-137 in Sea Water

Location	Sample volume analyzed (ℓ)	Cl (ℓ)	^{90}Sr		^{137}Cs	
			(mBq/ ℓ)	(mBq/ ℓ)	(mBq/ ℓ)	(mBq/ ℓ)
July, 1997						
Yoichi-bay, HOKKAIDO	40.0	18.72	1.6	\pm 0.27	2.5	\pm 0.30
Mutsu, AOMORI	60.0	18.60	1.7	\pm 0.30	3.4	\pm 0.37
Souma, FUKUSHIMA	40.0	17.11	1.4	\pm 0.26	2.2	\pm 0.28
Tokai, IBARAKI	40.0	17.23	1.5	\pm 0.31	1.9	\pm 0.29
Niigata, NIIGATA	46.0	17.74	1.9	\pm 0.29	2.7	\pm 0.30
Osaka-Port, OSAKA	40.0	9.88	1.9	\pm 0.29	1.6	\pm 0.28
August, 1997						
Mutsu, AOMORI	60.0	17.92	1.8	\pm 0.30	2.7	\pm 0.34
Ichihara, CHIBA	40.0	16.31	1.7	\pm 0.28	2.2	\pm 0.32
Yokosuka, KANAGAWA	40.0	17.97	1.7	\pm 0.28	2.1	\pm 0.27
Tokoname, AICHI	40.0	12.98	2.1	\pm 0.30	2.5	\pm 0.28
Yamaguchi-bay, YAMAGUCHI	40.0	17.11	1.6	\pm 0.28	2.7	\pm 0.33
Moji-Port, FUKUOKA	40.0	17.69	2.0	\pm 0.30	2.4	\pm 0.33
September, 1997						
Kaseda, KAGOSHIMA	40.0	----	2.0	\pm 0.27	2.2	\pm 0.27
Katsuren-machi, Okinawa	40.0	19.06	2.1	\pm 0.30	3.0	\pm 0.35

(7) Strontium-90 and Cesium-137 in Sea Sediments

(from Apr. 1997 to Sep. 1997)


-continued from No. 120 of this publication-

Table (7) Strontium-90 and Cesium-137 in Sea Sediments

Location	Depth (m)	^{90}Sr		^{137}Cs	
		(Bq/kg·dried Soil)	(Bq/kg·dried Soil)	(Bq/kg·dried Soil)	(Bq/kg·dried Soil)
July, 1997					
Yoichi-bay, HOKKAIDOU	13	0.044	\pm 0.059	0.53	\pm 0.091
Mutsu, AOMORI	15	0.10	\pm 0.091	0.25	\pm 0.071
Souma, FUKUSHIMA	5	0.082	\pm 0.051	0.52	\pm 0.91
Tokai-mura, IBARAKI	7	0.000	\pm 0.052	0.055	\pm 0.60
Niigata, NIIGATA	26	0.000	\pm 0.053	0.78	\pm 0.10
Osaka-Port, OSAKA	14.8	0.042	\pm 0.057	3.0	\pm 0.20
August, 1997					
Mutsu, AOMORI	15.0	0.17	\pm 0.079	4.6	\pm 0.25
Ichihara, CHIBA	15.0	0.14	\pm 0.070	2.6	\pm 0.18
Yokosuka, KANAGAWA	6	0.18	\pm 0.072	1.7	\pm 0.15
Tokoname, AICHI	22.0	0.14	\pm 0.066	3.6	\pm 0.21
Yamaguchi-bay, YAMAGUCHI	10	0.055	\pm 0.057	2.8	\pm 0.19
Moji-Port, FUKUOKA	10.0	0.026	\pm 0.052	1.1	\pm 0.13
September, 1997					
Kaseda, KAGOSHIMA	9	0.000	\pm 0.049	0.34	\pm 0.084
Katsuren-machi, Okinawa	11.0	0.090	\pm 0.081	0.029	\pm 0.076

* * Rain and Dry Fallout (for domestic program) * *

<Strontium-90>



<Cesium-137>





Fig. 1-1

* * Rain and Dry Fallout (for WHO program) * *

<Strontium-90>



<Cesium-137>





Fig.1—2

* * Air borne Dust * *

<Strontium-90>



<Cesium-137>





Fig. 2

* * Service Water (Source Water) * *

<Strontium-90>



<Cesium-137>





Fig. 3-1

* * Service Water (Tap Water) * *

<Strontium-90>



<Cesium-137>





Fig.3-2

* * Fresh Water * *

<Strontium-90>



<Cesium-137>





Fig. 4

* * Soil *

<Strontium-90>



<Cesium-137>





Fig. 5

* * Sea Water * *

<Strontium-90>



<Cesium-137>




Fig. 6

(38)
* * Sampling Locations in Japan * *

- | | |
|----------------|---------------|
| 1: Sapporo | 36: Hiroshima |
| 2: Aomori | 37: Kochi |
| 3: Morioka | 38: Matsuyama |
| 4: Akita | 39: Yamaguchi |
| 5: Sendai | 40: Ooita |
| 6: Yamagata | 41: Fukuoka |
| 7: Fukushima | 42: Saga |
| 8: Niigata | 43: Kumamoto |
| 9: Mito | 44: Miyazaki |
| 10: Utsunomiya | 45: Nagasaki |
| 11: Chiba | 46: Kagoshima |
| 12: Urawa | 47: Naha |
| 13: Shinjuku | |
| 14: Maebashi | |
| 15: Nagano | |
| 16: Yokohama | |
| 17: Toyama | |
| 18: Kouhu | |
| 19: Kanazawa | |
| 20: Shizuoka | |
| 21: Gifu | |
| 22: Fukui | |
| 23: Nagoya | |
| 24: Tsu | |
| 25: Ootsu | |
| 26: Kyoto | |
| 27: Nara | |
| 28: Osaka | |
| 29: Tottori | |
| 30: Kobe | |
| 31: Wakayama | |
| 32: Okayama | |
| 33: Matsue | |
| 34: Tokushima | |
| 35: Takamatsu | |

